

University of Galway Research Repository

Does bridging therapy in mechanical thrombectomy increase recanalization rates in ischemic stroke patients affected by acute large vessel occlusion?

Title	Does bridging therapy in mechanical thrombectomy increase recanalization rates in ischemic stroke patients affected by acute large vessel occlusion?
Author(s)	Silva Santos, Andreia M.;Rossi, Rosanna;Pandit, Abhay;Fitzgerald, Seán;Mereuta, Oana Madalina;Douglas, Andrew;Thornton, John;Brennan, Paul;Power, Sarah;O'Hare, Alan;Rentzos, Alexandros;Ceder, Erik;Tatlisumak, Turgut;Jood, Katarina;Redfors, Petra;Nordanstig, Annika;Duffy, Sharon;Gilvarry, Michael;McCarthy, Ray;Doyle, Karen M.
Publication Date	2019-05-22
Publication information	Silva Santos, Andreia M., Rossi, Rosanna, Pandit, Abhay, Fitzgerald, Seán, Mereuta, Oana Madalina, Douglas, Andrew, Thornton, John, Brennan, Paul, Power, Sarah, O'Hare, Alan, Rentzos, Alexandros, Ceder, Erik, Tatlisumak, Turgut, Jood, Katarina, Redfors, Petra, Nordanstig, Annika, Duffy, Sharon, Gilvarry, Michael, McCarthy, Ray, Doyle, Karen M. (2019). Does bridging therapy in mechanical thrombectomy increase recanalization rates in ischemic stroke patients affected by acute large vessel occlusion? Paper presented at the 5th European Stroke Organisation Conference (ESOC 2019), Milan, Italy, 22-24 May.
Publisher	SAGE Publications
Link to publisher's version	https://doi.org/10.1177/2396987319845581
Item record	http://hdl.handle.net/10379/15720

DOES BRIDGING-THERAPY IN MECHANICAL THROMBECTOMY INCREASE RECANALIZATION RATES IN ISCHEMIC STROKE PATIENTS AFFECTED BY ACUTE LARGE VESSEL OCCLUSION?

Andreia M. Silva Santos^{1,7*}, Rosanna Rossi^{1,2*}, Abhay Pandit², Seán Fitzgerald^{1,2}, Oana Madalina Mereuta^{1,2}, Andrew Douglas^{1,2}, John Thornton³, Paul Brennan³, Sarah Power³, Alan O' Hare³, Alexandros Rentzos⁴, Erik Ceder⁴, Turgut Tatlisumak⁵, Katarina Jood⁵, Petra Redfors⁵, Annika Nordanstig⁵, Sharon Duffy⁶, Michael Gilvarry⁶, Ray McCarthy⁶, Karen M. Doyle^{1,2}

*The two authors gave an equal contribution to the work.

¹Department of Physiology, School of Medicine, National University of Ireland, Galway, University Road, Galway, Ireland, ²CÚRAM—Centre for Research in Medical Devices, National University of Ireland Galway, Galway, Ireland, ³Department of Radiology, Royal college of Surgeons in Ireland, Beaumont Hospital, Dublin Ireland, ⁴Department of Interventional and Diagnostic Neuroradiology, Sahlgrenska University Hospital, Gothenburg, Sweden, ⁵Department of Neurology, Sahlgrenska University Hospital, Gothenburg, Sweden, ⁶Cerenovus, Galway, Ireland, ⁷Centro Universitário Unievangélica, Anápolis, Goiás, Brasil

Both intravenous thrombolysis with tissue plasminogen activator (IV-rtPA) and mechanical thrombectomy (MT) increase recanalization rates. We assessed if bridging-therapy (the concomitant use of rtPA and MT) could increase the recanalization rates and reduce the number of procedural passes in patients suffering from acute ischemic stroke (AIS) when compared to MT alone. Analysis of type of device used, stentriever or aspiration catheter, is also reported.

334 mechanically extracted thrombi were collected from two partner hospitals: Beaumont (Dublin) and Sahlgrenska (Gothenburg). 158 patients (47.3%) were treated with bridging-therapy, while 176 (52.7%) underwent MT alone. Recanalization rate was defined by using the modified Thrombolysis In Cerebral Infarction (mTICI) score. Non-parametric Kruskal-Wallis test was used for statistical analysis.

Bridging-therapy reduced the total number of passes to remove the clot (mean for MT+rtPA=2.27±2.10, MT alone=2.63±1.88, $H_1=4.376$, $p=0.036^*$). Analysing the device, rtPA lowered the overall number of passes using stentriever devices (mean for MT+rtPA=1.57±1.12, MT alone=2.36±1.48, $H_1=8.303$, $p=0.004^*$), but not for aspiration (mean for MT+rt-PA=1.78±1.22, MT alone=2.03±1.47, for $H_1=0.795$, $p=0.372$). Also, when using both devices no significant reduction of number of passes was observed (mean for MT+rtPA=3.29±2.90, MT alone=3.83±2.20, $H_1=3.027$, $p=0.082$). There was no significant effect on final mTICI score using bridging-therapy when compared to MT alone ($H_1=1.163$, $p=0.281$).

This small study suggests that bridging-therapy lowers the number of procedural passes in MT procedures, specifically when using stentriever devices. However, this did not have a significant effect on final mTICI score.

Funding: Science Foundation Ireland (Grant Number 13/RC/2073) and Cerenovus.