



Genus two partition and correlation functions for fermionic vertex operator superalgebras II

Title	Genus two partition and correlation functions for fermionic vertex operator superalgebras II
Author(s)	Tuite, Michael P.;Zuevsky, Alexander
Publication Date	2018-10-18

Genus Two Partition and Correlation Functions for Fermionic Vertex Operator Superalgebras II

Michael P. Tuite and Alexander Zuevsky*
School of Mathematics, Statistics and Applied Mathematics,
National University of Ireland Galway
University Road, Galway, Ireland

September 9, 2014

Abstract

We define and compute the continuous orbifold partition function and a generating function for all n -point correlation functions for the rank two free fermion vertex operator superalgebra on a genus two Riemann surface formed by self-sewing a torus. The partition function is proportional to an infinite dimensional determinant with entries arising from torus Szegő kernel and the generating function is proportional to a finite determinant of genus two Szegő kernels. These results follow from an explicit analysis of all torus n -point correlation functions for intertwiners of the irreducible modules of the Heisenberg vertex operator algebra. We prove that the partition and n -point correlation functions are holomorphic on a suitable domain and describe their modular properties. We also describe an identity for the genus two Riemann theta series analogous to the Jacobi triple product identity.

*Supported by a Science Foundation Ireland Research Frontiers Programme Grant, and by Max-Planck Institut für Mathematik, Bonn

1 Introduction

This paper is a continuation of the study in the theory of Vertex Operator Algebras (VOAs) [FLM, FHL, K, MN, MT5] of partition functions and correlation functions on a genus two Riemann surface [T, MT1, MT2, MT3, MT4, TZ2]. Such a surface can be constructed by sewing two separate tori together, which we refer to as the ϵ -formalism, or by self-sewing a single torus, which we refer to as the ρ -formalism [MT2]. In our approach, we define the genus two partition and n -point correlation functions in terms of genus one VOA data combined in accordance with the given sewing scheme. This is based solely on the properties of the VOA with no assumed analytic or modular properties for partition or correlation functions. This is in contrast to other approaches to correlation functions using algebraic geometric methods in mathematics (e.g. [TUY, Z2]) or other approaches taken in physics (e.g. [EO, DP, DVPFHLS]).

In the companion paper ref. [TZ2] we studied the continuous orbifold partition and correlation functions for the rank two free fermion Vertex Operator Superalgebra (VOSA) $V(H, \mathbb{Z} + \frac{1}{2})^{\otimes 2}$ on a genus two surface in the ϵ -formalism. The present paper studies this system in the ρ -formalism where a genus two Riemann surface is constructed by sewing a handle at two points, separated by w , to a torus with modular parameter τ and with sewing parameter ρ giving a family of surfaces parametrized by $(\tau, w, \rho) \in \mathcal{D}^\rho$, a particular sewing domain. Although there are some features in common with ref. [TZ2], the present ρ -formalism approach is more subtle in two important respects: firstly, we must consider genus one correlation functions for intertwining vertex operators associated with g -twisted modules for a continuous winding automorphism g and secondly, the genus two partition and n -point functions obtained are multi-valued on \mathcal{D}^ρ . There is no Zhu reduction theory available for general genus one intertwiner correlation functions unlike the reduction theory developed in [MTZ] and used in the ϵ -formalism. Instead, we use bosonization and compute all torus n -point functions for the Generalized VOA $\mathcal{M} = \bigoplus_{\alpha \in \mathbb{C}} M \otimes e^\alpha$ formed by the intertwiners for the rank one Heisenberg VOA M and all of its irreducible modules $M \otimes e^\alpha$ for all $\alpha \in \mathbb{C}$ [BK, TZ3]. This requires an extension of previous results for genus one lattice theory correlation functions to the Generalized VOA \mathcal{M} . We also make repeated use of an explicit expression of genus two Szegő kernel in terms of genus one data in the ρ -formalism described in [TZ1]. The genus two partition and n -point functions are computed explicitly and involve an infinite

determinant whose components arise from genus one Szegő kernel data. The genus two partition and n -point functions are multi-valued functions on \mathcal{D}^ρ that lift to holomorphic functions on a suitable covering space and are modular invariant under a subgroup of the genus two symplectic group $\mathrm{Sp}(4, \mathbb{Z})$. We also compute the partition function in a bosonic basis and obtain an analogue of the classic Jacobi triple product formula for the genus two Riemann theta series.

Section 2 contains a review of the ρ -formalism for the construction of a genus two Szegő kernel $S^{(2)}$ on \mathcal{D}^ρ in terms of elliptic Szegő kernel data and given multipliers on the four cycles of a genus two Riemann surface formed by self-sewing a torus [MT2, TZ1]. We introduce a certain infinite matrix T constructed from the genus one Szegő data and the multipliers which plays a central role: $(I-T)^{-1}$ appears explicitly in the formula for $S^{(2)}$ and $\det(I-T)$, which is holomorphic on \mathcal{D}^ρ , appears in our later VOA results.

Section 3 reviews the rank two free fermion Vertex Operator Superalgebra (VOA) $V(H, \mathbb{Z} + \frac{1}{2})^{\otimes 2}$ and its twisted modules for a continuous winding automorphism $g = e^{-2\pi i \alpha a(0)}$ for Heisenberg vector a . We often make use of bosonization whereby $V(H, \mathbb{Z} + \frac{1}{2})^{\otimes 2} \cong V_{\mathbb{Z}}$, the \mathbb{Z} -lattice VOA with trivial cocycle structure. In particular, we consider the interwining operators on $\mathcal{M} = \bigoplus_{\alpha \in \mathbb{C}} M \otimes e^\alpha$ that satisfy a Generalized VOA described in detail in [TZ3, BK]. This leads to generalized notions of locality, skew-symmetry, associativity, commutivity with respect to the Heisenberg sub VOA and an invariant linear form on \mathcal{M} .

In Section 4 we compute general n -point intertwiner correlation functions for elements of \mathcal{M} in terms of elliptic prime forms and elliptic and quasi-modular forms. These expressions are natural generalizations of results for lattice VOAs [MT1] extended to \mathcal{M} . This allows us to compute a generating form for all intertwiner torus n -point functions for $V_{\mathbb{Z}}$ expressed in terms of a finite determinant of genus one Szegő kernels.

Section 5 contains our main results concerning the genus two partition function and a generating form for all genus two n -point functions. These are defined in terms of four commuting automorphisms generated by $a(0)$ which determine the $S^{(2)}$ multipliers. The partition function which is defined as a sum over a basis for a g -twisted module of particular genus one interwining 2-point functions is explicitly computed in Theorem 5.1. The result is proportional to $\det(I-T)$ but is multivalued on \mathcal{D}^ρ due to a separate factor

of $(-\rho/K(w, \tau)^2)^{\frac{1}{2}\kappa^2}$ where $g = e^{-2\pi i \kappa a(0)}$ for $\kappa \in (-\frac{1}{2}, \frac{1}{2})$ and $K(w, \tau)$ is the elliptic prime form. In Theorem 5.10 we show that the partition function lifts to a holomorphic function on a covering space $\widehat{\mathcal{D}}^\rho$ and prove that it is modular invariant under a particular subgroup $L \subset \mathrm{Sp}(4, \mathbb{Z})$, the genus two symplectic group. We also compute the partition function in an alternative bosonic basis and obtain an analogue of the Jacobi triple product formula for the genus two Riemann theta series in Theorem 5.9. Lastly, in Theorem 5.3 a modular invariant generating form for all genus two n -point correlation functions is computed in terms of the genus two partition function and a finite determinant of genus two Szegő kernels.

2 The Szegő Kernel in the ρ -Formalism

In this section we consider the construction of a genus two Riemann surface formed by self-sewing a handle to a torus using an explicit sewing scheme which we refer to as the ρ -formalism. In particular, we review the construction of an explicit formula for the genus two Szegő kernel in terms of Szegő kernel on the torus. Further details can be found in [MT2] and [TZ1].

2.1 The Szegő Kernel on a Riemann Surface

Consider a compact connected Riemann surface Σ of genus g with canonical homology cycle basis a_i, b_i for $i = 1, \dots, g$. Let ν_i be a basis of holomorphic 1-forms with normalization $\oint_{a_i} \nu_j = 2\pi i \delta_{ij}$ and period matrix $\Omega_{ij} = \frac{1}{2\pi i} \oint_{b_i} \nu_j \in \mathbb{H}_g$, the Siegel upper half plane. Define the theta function with real characteristics [M, F, FK]

$$\vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} (z|\Omega) = \sum_{n \in \mathbb{Z}^g} e^{i\pi(n+\alpha) \cdot \Omega \cdot (n+\alpha) + (n+\alpha) \cdot (z+2\pi i \beta)}, \quad (1)$$

for $\alpha = (\alpha_j), \beta = (\beta_j) \in \mathbb{R}^g$ and $z = (z_j) \in \mathbb{C}^g$ for $j = 1, \dots, g$. The Szegő kernel is defined for $\vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} (0|\Omega) \neq 0$ by [S, HS, F]

$$S \begin{bmatrix} \theta \\ \phi \end{bmatrix} (x, y|\Omega) = \frac{\vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \left(\int_y^x \nu \mid \Omega \right)}{\vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} (0|\Omega) E(x, y)}, \quad (2)$$

for $x, y \in \Sigma$ and where $\theta = (\theta_j)$, $\phi = (\phi_j) \in U(1)^n$ for

$$\theta_j = -e^{-2\pi i\beta_j}, \quad \phi_j = -e^{2\pi i\alpha_j}, \quad j = 1, \dots, g. \quad (3)$$

(where the -1 factors are included for later convenience) and $E(x, y)$ is the prime form [M, F]. The Szegő kernel is periodic in x along the a_i and b_j cycles with multipliers $-\phi_i$ and $-\theta_j$ respectively and is a meromorphic $(\frac{1}{2}, \frac{1}{2})$ -form (and is thus necessarily defined on a double-cover $\tilde{\Sigma}$ of the Riemann surface) satisfying¹

$$\begin{aligned} S \begin{bmatrix} \theta \\ \phi \end{bmatrix} (x, y) &\sim \frac{1}{x-y} dx^{\frac{1}{2}} dy^{\frac{1}{2}} \quad \text{for } x \sim y, \\ S \begin{bmatrix} \theta \\ \phi \end{bmatrix} (x, y) &= -S \begin{bmatrix} \theta^{-1} \\ \phi^{-1} \end{bmatrix} (y, x), \end{aligned}$$

where $\theta^{-1} = (\theta_i^{-1})$ and $\phi^{-1} = (\phi_i^{-1})$. The Szegő kernel transforms under the action of the symplectic modular group for $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \text{Sp}(2g, \mathbb{Z})$ as follows [F]

$$S \begin{bmatrix} \tilde{\theta} \\ \tilde{\phi} \end{bmatrix} (x, y | \tilde{\Omega}) = S \begin{bmatrix} \theta \\ \phi \end{bmatrix} (x, y | \Omega), \quad (4)$$

$$\tilde{\Omega} = (A\Omega + B)(C\Omega + D)^{-1}, \quad (5)$$

where $\tilde{\theta}_j = -e^{-2\pi i\tilde{\beta}_j}$, $\tilde{\phi}_j = -e^{2\pi i\tilde{\alpha}_j}$ with

$$\begin{bmatrix} -\tilde{\beta} \\ \tilde{\alpha} \end{bmatrix} = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} -\beta \\ \alpha \end{bmatrix} + \frac{1}{2} \begin{bmatrix} -\text{diag}(AB^T) \\ \text{diag}(CD^T) \end{bmatrix}, \quad (6)$$

where $\text{diag}(M)$ denotes the diagonal elements of a matrix M .

2.2 Genus Two Riemann Surface Formed from Self-Sewing a Torus

We now briefly review the construction of a genus two Riemann surface $\Sigma^{(2)}$ formed by self-sewing a handle to a torus which we refer to this as the ρ -formalism [MT2, TZ1]. Let $\Sigma^{(1)} = \mathbb{C}/\Lambda$ denote an oriented torus for lattice $\Lambda = 2\pi i(\mathbb{Z}\tau \oplus \mathbb{Z})$ and $\tau \in \mathbb{H}_1$. Define annuli \mathcal{A}_a , $a = 1, 2$ centered at $z = 0$

¹We use the convention $E(x, y) \sim (x-y)dx^{-\frac{1}{2}}dy^{-\frac{1}{2}}$ for $x \sim y$.

and $z = w$ of $\Sigma^{(1)}$ with local coordinates $z_1 = z$ and $z_2 = z - w$ respectively. Define the following label convention

$$\bar{1} = 2, \quad \bar{2} = 1. \quad (7)$$

Take the outer radius of \mathcal{A}_a to be $r_a < \frac{1}{2}D(q)$ for $D(q) = \min_{\lambda \in \Lambda, \lambda \neq 0} |\lambda|$ and the inner radius to be $|\rho|/r_{\bar{a}}$, with $|\rho| \leq r_1 r_2$, where we introduce a complex parameter ρ , $|\rho| \leq r_1 r_2$. r_1, r_2 must be sufficiently small to ensure that the disks do not intersect. Excise the disks

$$\{z_a, |z_a| < |\rho| r_{\bar{a}}^{-1}\} \subset \Sigma^{(1)},$$

to form a twice-punctured surface

$$\widehat{\Sigma}^{(1)} = \Sigma^{(1)} \setminus \bigcup_{a=1,2} \{z_a, |z_a| < |\rho| r_{\bar{a}}^{-1}\}.$$

We define annular regions $\mathcal{A}_a \subset \widehat{\Sigma}^{(1)}$ with $\mathcal{A}_a = \{z_a, |\rho| r_{\bar{a}}^{-1} \leq |z_a| \leq r_a\}$ and identify them as a single region $\mathcal{A} = \mathcal{A}_1 \simeq \mathcal{A}_2$ via the sewing relation

$$z_1 z_2 = \rho, \quad (8)$$

to form a compact genus two Riemann surface $\Sigma^{(2)} = \widehat{\Sigma}^{(1)} \setminus \{\mathcal{A}_1 \cup \mathcal{A}_2\} \cup \mathcal{A}$, parameterized by [MT2]

$$\mathcal{D}^\rho = \{(\tau, w, \rho) \in \mathbb{H}_1 \times \mathbb{C} \times \mathbb{C}, |w - \lambda| > 2|\rho|^{\frac{1}{2}} > 0, \lambda \in \Lambda\}. \quad (9)$$

2.3 The Genus Two Szegő Kernel in the ρ -Formalism

Now we recall the construction of the Szegő kernel on a genus two Riemann surface in the ρ -formalism [TZ1]. The genus one prime form for $x, y \in \mathbb{C}$ and $\tau \in \mathbb{H}_1$ is

$$E(x, y) = K(x - y, \tau) dx^{-\frac{1}{2}} dy^{-\frac{1}{2}}, \quad (10)$$

where

$$K(z, \tau) = \frac{\vartheta_1(z, \tau)}{\partial_z \vartheta_1(0, \tau)}, \quad \vartheta_1(z, \tau) = \vartheta \left[\begin{matrix} \frac{1}{2} \\ \frac{1}{2} \end{matrix} \right] (z, \tau). \quad (11)$$

Let $(\theta_1, \phi_1) \in U(1) \times U(1)$ with $(\theta_1, \phi_1) \neq (1, 1)$. The genus one Szegő kernel is

$$S^{(1)} \left[\begin{matrix} \theta_1 \\ \phi_1 \end{matrix} \right] (x, y | \tau) = P_1 \left[\begin{matrix} \theta_1 \\ \phi_1 \end{matrix} \right] (x - y, \tau) dx^{\frac{1}{2}} dy^{\frac{1}{2}}, \quad (12)$$

where

$$P_1 \begin{bmatrix} \theta_1 \\ \phi_1 \end{bmatrix} (z, \tau) = \frac{\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (z, \tau)}{\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (0, \tau)} \frac{1}{K(z, \tau)}, \quad (13)$$

where $-\phi_1 = \exp(2\pi i \alpha_1)$ and $-\theta_1 = \exp(-2\pi i \beta_1)$ are the periodicities of $S^{(1)} \begin{bmatrix} \theta_1 \\ \phi_1 \end{bmatrix} (x, y | \tau)$ in x on the standard a and b cycles respectively.

In [TZ1] we determine the genus two Szegő kernel

$$S^{(2)}(x, y) = S^{(2)} \begin{bmatrix} \theta^{(2)} \\ \phi^{(2)} \end{bmatrix} (x, y),$$

with periodicities $(\theta^{(2)}, \phi^{(2)}) = (\theta_i, \phi_i)$ for $i = 1, 2$ on a natural homology basis on the genus two Riemann surface $\Sigma^{(2)}$ formed by self-sewing the torus $\Sigma^{(1)}$ in terms of genus one Szegő kernel data $S^{(1)}(x, y) = S^{(1)} \begin{bmatrix} \theta_1 \\ \phi_1 \end{bmatrix} (x, y)$ with $(\theta_1, \phi_1) \neq (1, 1)$. The $S^{(2)}$ multipliers (3) on the cycles a_1, b_1 are determined by the corresponding multipliers of $S^{(1)}$ so that $\phi_1^{(2)} = \phi_1$ and $\theta_1^{(2)} = \theta_1$ i.e. $\alpha_1^{(2)} = \alpha_1$ and $\beta_1^{(2)} = \beta_1$. The remaining multipliers $\phi_2 = \phi_2^{(2)} = -e^{2\pi i \alpha_2^{(2)}}$ and $\theta_2 = \theta_2^{(2)} = -e^{-2\pi i \beta_2^{(2)}}$ on the cycles a_2 and b_2 must be specified by the additional conditions

$$\begin{aligned} S^{(2)}(e^{2\pi i} x_a, y) &= -\phi_2^{a-\bar{a}} S^{(2)}(x_a, y), \\ S^{(2)}(x_a, y) &= -\theta_2^{a-\bar{a}} S^{(2)}(x_{\bar{a}}, y), \end{aligned}$$

for $x_a \in \mathcal{A}_a$ and $x_{\bar{a}} \in \mathcal{A}_{\bar{a}}$ (recalling the label convention (7)).

It is convenient to define $\kappa \in (\frac{1}{2}, \frac{1}{2})$ by $\phi_2 = -e^{2\pi i \kappa}$ i.e. $\kappa = \alpha_2^{(2)} \pmod{1}$. We then find that $S^{(2)}$ is holomorphic in ρ for $|\rho| < r_1 r_2$ with [TZ1]

$$S^{(2)}(x, y) = S_\kappa(x, y) + O(\rho),$$

for $x, y \in \widehat{\Sigma}^{(1)}$ and

$$S_\kappa(x, y) = \left(\frac{\vartheta_1(x-w, \tau) \vartheta_1(y, \tau)}{\vartheta_1(x, \tau) \vartheta_1(y-w, \tau)} \right)^\kappa \frac{\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (x-y + \kappa w, \tau)}{\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (\kappa w, \tau) K(x-y, \tau)} dx^{\frac{1}{2}} dy^{\frac{1}{2}}, \quad (14)$$

for $\kappa \neq -\frac{1}{2}$ (with a different expression when $\kappa = -\frac{1}{2}$ given in [TZ1]). We will assume $\kappa \neq -\frac{1}{2}$ throughout this paper. Note also that $S_{\kappa=0}(x, y) = S^{(1)} \left[\begin{smallmatrix} \theta_1 \\ \phi_1 \end{smallmatrix} \right] (x, y)$, the genus one Szegő kernel.

$S_\kappa(x, y)$ has an expansion in the neighborhood of the punctures at 0, w in terms of local coordinates $x_1 = x, y_1 = y$ and $x_1 = x - w, y_2 = y - w$ as follows [TZ3]

$$S_\kappa(x_{\bar{a}}, y_b) = \left[\delta_{\bar{a},b} \frac{1}{x_b - y_b} \left(\frac{x_b}{y_b} \right)^{\kappa(-1)^b} + \sum_{k,l \geq 1} C_{ab}(k, l) x_{\bar{a}}^{k_a-1} y_b^{l_b-1} \right] dx_{\bar{a}}^{\frac{1}{2}} dy_b^{\frac{1}{2}}, \quad (15)$$

where $C_{ab}(k, l) = C_{ab} \left[\begin{smallmatrix} \theta_1 \\ \phi_1 \end{smallmatrix} \right] (k, l | \tau, w, \kappa)$ and $k_a = k + \kappa(-1)^{\bar{a}}$ for integer $k \geq 1$ and $a = 1, 2$. We may invert this to obtain the infinite block moment matrix

$$C_{ab}(k, l) = \frac{1}{(2\pi i)^2} \oint_{\mathcal{C}_{\bar{a}}(x_{\bar{a}})} \oint_{\mathcal{C}_b(y_b)} (x_{\bar{a}})^{-k_a} (y_b)^{-l_b} S_\kappa(x_{\bar{a}}, y_b) dx_{\bar{a}}^{\frac{1}{2}} dy_b^{\frac{1}{2}}. \quad (16)$$

We also define half-order differentials

$$d_a(x, k) = \frac{1}{2\pi i} \oint_{\mathcal{C}_a(y_a)} y_a^{-k_a} S_\kappa(x, y_a) dy_a^{\frac{1}{2}}, \quad (17)$$

$$\bar{d}_a(y, k) = \frac{1}{2\pi i} \oint_{\mathcal{C}_{\bar{a}}(x_{\bar{a}})} x_{\bar{a}}^{-k_a} S_\kappa(x_{\bar{a}}, y) dx_{\bar{a}}^{\frac{1}{2}}. \quad (18)$$

We further define

$$G_{ab}(k, l) = \rho^{\frac{1}{2}(k_a+l_b-1)} C_{ab}(k, l), \quad (19)$$

$$h_a(x, k) = \rho^{\frac{1}{2}(k_a-\frac{1}{2})} d_a(x, k), \quad (20)$$

$$\bar{h}_a(y, k) = \rho^{\frac{1}{2}(k_a-\frac{1}{2})} \bar{d}_a(y, k), \quad (21)$$

with associated infinite matrix $G = (G_{ab}(k, l))$ and row vectors $h(x) = (h_a(x, k))$ and $\bar{h}(y) = (\bar{h}_a(y, k))$. From the sewing relation (8) we have

$$dz_{\bar{a}}^{\frac{1}{2}} = (-1)^{\bar{a}} \xi \rho^{\frac{1}{2}} \frac{dz_{\bar{a}}^{\frac{1}{2}}}{z_{\bar{a}}}, \quad (22)$$

for $\xi \in \{\pm\sqrt{-1}\}$ depending on the branch of the double cover of $\Sigma^{(1)}$ chosen. Finally define

$$T = \xi G D^{\theta_2}, \quad (23)$$

with an infinite diagonal matrix

$$D^{\theta_2}(k, l) = \begin{bmatrix} \theta_2^{-1} & 0 \\ 0 & -\theta_2 \end{bmatrix} \delta(k, l). \quad (24)$$

Defining $\det(I - T)$ by $\log \det(I - T) = \text{Tr} \log(I - T) = -\sum_{n \geq 1} \frac{1}{n} \text{Tr} T^n$, we find

Theorem 2.1 ([TZ1]) *On the ρ -sewing domain \mathcal{D}^ρ we have:*

- (i) $(I - T)^{-1} = \sum_{n \geq 0} T^n$ is convergent.
- (ii) $\det(I - T)$ is holomorphic and non-vanishing.
- (iii) The genus two Szegő kernel for $x, y \in \widehat{\Sigma}^{(1)}$ is given by

$$S^{(2)}(x, y) = S_\kappa(x, y) + \xi h(x) D^\theta (I - T)^{-1} \bar{h}^T(y), \quad (25)$$

where $\bar{h}^T(y)$ denotes the transpose of the infinite row vector $\bar{h}(y)$.

3 The Free Fermion VOSA and its Twisted Modules

3.1 The Free Fermion VOSA

We consider in this paper the rank two free fermion Vertex Operator Super Algebra (VOSA) $V = V(H, \mathbb{Z} + \frac{1}{2})^{\otimes 2}$ of central charge 1 (e.g. see [K, MTZ] for details). The weight $\frac{1}{2}$ space is spanned by ψ^+, ψ^- with vertex operator modes which satisfy the anti-commutation relations

$$[\psi^+(m), \psi^-(n)] = \delta_{m, -n-1}, \quad [\psi^+(m), \psi^+(n)] = [\psi^-(m), \psi^-(n)] = 0.$$

V is spanned by Fock vectors of the form

$$\Psi(\mathbf{k}, \mathbf{l}) \equiv \psi^+(-k_1) \dots \psi^+(-k_s) \psi^-(-l_1) \dots \psi^-(-l_t) \mathbf{1}, \quad (26)$$

for distinct $0 < k_1 < \dots < k_s$ and $0 < l_1 < \dots < l_t$, and of Virasoro weight

$$\text{wt}(\Psi(\mathbf{k}, \mathbf{l})) = \sum_{i=1}^s \left(k_i - \frac{1}{2} \right) + \sum_{j=1}^t \left(l_j - \frac{1}{2} \right), \quad (27)$$

with Virasoro vector

$$\omega = \frac{1}{2}(\psi^+(-2)\psi^-(-1) + \psi^-(-2)\psi^+(-1))\mathbf{1}.$$

The weight 1 space is spanned by $a = \psi^+(-1)\psi^-$ whose modes obey the Heisenberg commutation relations

$$[a(m), a(n)] = m\delta_{m,-n}.$$

As is well known, we may decompose V into irreducible M -modules $M \otimes e^m$ with $a(0)$ eigenvalue $m \in \mathbb{Z}$ so that $V = V(H, \mathbb{Z} + \frac{1}{2})^{\otimes 2} \cong V_{\mathbb{Z}} = \bigoplus_{m \in \mathbb{Z}} M \otimes e^m$, the lattice VOSA for the \mathbb{Z} -lattice with trivial cocycle structure.

3.2 g -Twisted $V_{\mathbb{Z}}$ -Modules and a Generalized VOA

$a(0)$ generates continuous winding $V_{\mathbb{Z}}$ -automorphism $g = e^{-2\pi i \alpha a(0)}$ for $\alpha \in \mathbb{C}$. In particular, the fermion number involution is $\sigma = e^{\pi i a(0)}$. We define for all $u \in M$ the following operators

$$Y_g(u \otimes e^m, z) = Y(\Delta(\alpha, z)(u \otimes e^m), z), \quad (28)$$

$$\Delta(\alpha, z) = z^{\alpha a(0)} Y_+(\alpha, -z), \quad (29)$$

$$Y_{\pm}(\alpha, z) = \exp\left(\mp \alpha \sum_{n>0} \frac{a(\pm n)}{n} z^{\mp n}\right). \quad (30)$$

Then we have [Li]

Proposition 3.1 $(V_{\mathbb{Z}}, Y_g)$ is a g -twisted $V_{\mathbb{Z}}$ -module.

In Section 5 of [TZ3] an isomorphic construction is described whereby the g -twisted module is determined by the action of the original vertex operators on a twisted vector space $V_{\mathbb{Z}+\alpha} = e^{\alpha \hat{q}} V_{\mathbb{Z}} = \bigoplus_{m \in \mathbb{Z}} M \otimes e^{m+\alpha}$ where

$$Y_g(u \otimes e^m, z) = e^{-\alpha \hat{q}} Y(u \otimes e^m, z) e^{\alpha \hat{q}}, \quad (31)$$

where \hat{q} is defined by

$$[a(n), \hat{q}] = \delta_{n,0}. \quad (32)$$

In particular $Y_g(\omega, z)$ determines the g -twisted grading operator

$$L_g(0) = L(0) + \alpha a(0) + \frac{1}{2} \alpha^2. \quad (33)$$

Hence $\Psi(\mathbf{k}, \mathbf{l}) \in M \otimes e^{s-t}$ of (26) has g -twisted Virasoro weight $\text{wt}(\Psi(\mathbf{k}, \mathbf{l})) + \alpha(s-t) + \frac{1}{2}\alpha^2$ which is equal to the $L(0)$ weight of the $V_{\mathbb{Z}+\alpha}$ twisted Fock vector

$$\Psi_\alpha(\mathbf{k}, \mathbf{l}) \equiv e^{\alpha\hat{q}}\Psi(\mathbf{k}, \mathbf{l}) \in M \otimes e^{s-t+\alpha}. \quad (34)$$

In [TZ3] we describe a Generalized VOA with vector space

$$\mathcal{M} = \bigoplus_{\alpha \in \mathbb{C}} M \otimes e^\alpha, \quad (35)$$

formed as a direct sum of the Heisenberg VOA with all of its irreducible modules. \mathcal{M} is spanned by $\Psi_\alpha(\mathbf{k}, \mathbf{l})$ for all $\alpha \in \mathbb{C}$. The generalized vertex operators are

$$\mathcal{Y}(u \otimes e^\alpha, z) = e^{\alpha\hat{q}} Y_-(\alpha, z) Y(u, z) Y_+(\alpha, z) z^{\alpha a(0)}. \quad (36)$$

(36) reduces to the usual bosonized form of the vertex operators for $V_{\mathbb{Z}}$ for $\alpha \in \mathbb{Z}$. A similar construction also appears in [BK]. The vertex operators (36) possess the creativity and translation property with Heisenberg Virasoro vector and satisfy a generalized Jacobi identity (with a trivial cocycle structure):

Theorem 3.2 *($\mathcal{M}, \mathcal{Y}, \mathbf{1}, \omega$) is a Generalized VOA with vertex operators $\mathcal{Y}(u \otimes e^\alpha, z) \in \text{End}(\mathcal{M})\{z\}$ (the space of $\text{End}(\mathcal{M})$ -valued formal series in z with arbitrary complex powers of z) obeying the generalized Jacobi identity*

$$\begin{aligned} & z_0^{-1} \left(\frac{z_1 - z_2}{z_0} \right)^{-\alpha\beta} \delta \left(\frac{z_1 - z_2}{z_0} \right) \mathcal{Y}(u \otimes e^\alpha, z_1) \mathcal{Y}(v \otimes e^\beta, z_2) \\ & - z_0^{-1} \left(\frac{z_2 - z_1}{z_0} \right)^{-\alpha\beta} \delta \left(\frac{z_2 - z_1}{-z_0} \right) \mathcal{Y}(v \otimes e^\beta, z_2) \mathcal{Y}(u \otimes e^\alpha, z_1) \\ & = z_2^{-1} \delta \left(\frac{z_1 - z_0}{z_2} \right) \mathcal{Y}(\mathcal{Y}(u \otimes e^\alpha, z_0)(v \otimes e^\beta), z_2) \left(\frac{z_1 - z_0}{z_2} \right)^{\alpha a(0)}, \end{aligned} \quad (37)$$

for all $u \otimes e^\alpha, v \otimes e^\beta \in \mathcal{M}$.

The Generalized VOA leads to more general notions of locality, skew-symmetry, associativity and commutivity than those for a VOSA as follows:

Proposition 3.3 For $u \otimes e^\alpha, v \otimes e^\beta, w \otimes e^\gamma \in \mathcal{M}$ and for integer $N \gg 0$

$$\begin{aligned} & (z_1 - z_2)^N (z_1 - z_2)^{-\alpha\beta} \mathcal{Y}(u \otimes e^\alpha, z_1) \mathcal{Y}(v \otimes e^\beta, z_2) \\ &= (z_1 - z_2)^N (z_2 - z_1)^{-\alpha\beta} \mathcal{Y}(v \otimes e^\beta, z_2) \mathcal{Y}(u \otimes e^\alpha, z_1), \end{aligned} \quad (38)$$

$$z^{-\alpha\beta} \mathcal{Y}(u \otimes e^\alpha, z) v \otimes e^\beta = (-z)^{-\alpha\beta} e^{zL(-1)} \mathcal{Y}(v \otimes e^\beta, -z) u \otimes e^\alpha, \quad (39)$$

$$\begin{aligned} & (z_0 + z_2)^{N-\alpha\gamma} \mathcal{Y}(u \otimes e^\alpha, z_0 + z_2) \mathcal{Y}(v \otimes e^\beta, z_2) w \otimes e^\gamma \\ &= (z_2 + z_0)^{N-\alpha\gamma} \mathcal{Y}(\mathcal{Y}(u \otimes e^\alpha, z_0) (v \otimes e^\beta), z_2) w \otimes e^\gamma, \end{aligned} \quad (40)$$

$$[u(k), \mathcal{Y}(v \otimes e^\beta, z)] = \sum_{j \geq 0} \binom{k}{j} \mathcal{Y}(u(j) v \otimes e^\beta, z) z^{k-j}, \quad u \in M. \quad (41)$$

The proof appears in Appendix 6.1.

It is convenient to define for formal parameter z and $\chi \in \mathbb{C}$

$$(-z)^\chi = e^{i\pi B \chi} z^\chi, \quad (42)$$

where we choose, once and for all, an *odd integer* B parametrizing the formal branch cut². Then generalized locality and skew-symmetry can be rewritten as

$$\begin{aligned} & (z_1 - z_2)^{N-\alpha\beta} \mathcal{Y}(u \otimes e^\alpha, z_1) \mathcal{Y}(v \otimes e^\beta, z_2) \\ &= e^{-i\pi B \alpha\beta} (z_1 - z_2)^{N-\alpha\beta} \mathcal{Y}(v \otimes e^\beta, z_2) \mathcal{Y}(u \otimes e^\alpha, z_1), \end{aligned} \quad (43)$$

$$\mathcal{Y}(u \otimes e^\alpha, z) v \otimes e^\beta = e^{-i\pi B \alpha\beta} e^{zL(-1)} \mathcal{Y}(v \otimes e^\beta, -z) u \otimes e^\alpha. \quad (44)$$

3.2.1 An Invariant Form on \mathcal{M}

In [TZ3] we introduced an invariant bilinear form $\langle \cdot, \cdot \rangle$ on \mathcal{M} associated with the Möbius map [FHL], [S], [TZ2]

$$\begin{pmatrix} 0 & \lambda \\ -e^{i\pi B} \lambda^{-1} & 0 \end{pmatrix} : z \mapsto -\frac{\lambda^2}{e^{i\pi B} z}, \quad (45)$$

for $\lambda \neq 0$. We are particularly interested in the Möbius map $z \mapsto \rho/z$ associated with the sewing condition (8) so that we will later choose

$$\lambda = e^{\frac{1}{2}i\pi B} \rho^{\frac{1}{2}}, \quad (46)$$

²Note some notational changes from [TZ3]

for the odd integer B of (42). Thus we reformulate the sewing relationship (8) as $z_1 = -\frac{\lambda^2}{z_2}$ so that (22) implies $dz_1^{\frac{1}{2}} = \xi \rho^{\frac{1}{2}} / z_2 dz_2^{\frac{1}{2}}$ for $\xi = e^{\frac{1}{2}i\pi B}$.

Define the adjoint vertex operator

$$\mathcal{Y}^\dagger(u \otimes e^\alpha, z) = \mathcal{Y}\left(e^{-z\lambda^{-2}L(1)}\left(\frac{\lambda}{e^{i\pi B}z}\right)^{2L(0)}(u \otimes e^\alpha), \frac{\lambda^2}{e^{i\pi B}z}\right). \quad (47)$$

A bilinear form $\langle \cdot, \cdot \rangle_\lambda$ on \mathcal{M} is said to be invariant if for all $u \otimes e^\alpha, v \otimes e^\beta, w \otimes e^\gamma \in \mathcal{M}$ we have

$$\langle \mathcal{Y}(u \otimes e^\alpha, z)(v \otimes e^\beta), w \otimes e^\gamma \rangle_\lambda = e^{-i\pi B\alpha\beta} \langle v \otimes e^\beta, \mathcal{Y}^\dagger(u \otimes e^\alpha, z)w \otimes e^\gamma \rangle_\lambda. \quad (48)$$

(47) reduces to the usual definition for a VOSA when $\alpha, \beta, \gamma \in \mathbb{Z}$ [S, TZ2]. Choosing the normalization $\langle \mathbf{1}, \mathbf{1} \rangle = 1$ then $\langle \cdot, \cdot \rangle_\lambda$ on \mathcal{M} is symmetric, unique and invertible with [TZ3]

$$\langle u \otimes e^\alpha, v \otimes e^\beta \rangle_\lambda = \lambda^{-\alpha^2} \delta_{\alpha, -\beta} \langle u \otimes e^0, v \otimes e^0 \rangle_\lambda. \quad (49)$$

The dual of the Fock vector $\Psi = \Psi(\mathbf{k}, \mathbf{l})$ with respect to $\langle \cdot, \cdot \rangle_\lambda$, which we refer to as the λ -dual, is

$$\bar{\Psi}(\mathbf{k}, \mathbf{l}) = (-1)^{st + \lfloor \text{wt}(\Psi) \rfloor} \lambda^{2\text{wt}(\Psi)} \Psi(\mathbf{l}, \mathbf{k}),$$

where $\lfloor x \rfloor$ denotes the integer part of x [TZ2]. Applying (46) and (49) it follows that $\bar{\Psi}_\alpha = \bar{\Psi}_\alpha(\mathbf{k}, \mathbf{l})$ of (34) has λ -dual

$$\begin{aligned} \bar{\Psi}_\alpha(\mathbf{k}, \mathbf{l}) &= (-1)^{st + \lfloor \text{wt}(\Psi_\alpha) \rfloor} \lambda^{2\text{wt}(\Psi_\alpha)} \Psi_{-\alpha}(\mathbf{l}, \mathbf{k}) \\ &= (-1)^{st + \lfloor \text{wt}(\Psi_\alpha) \rfloor} e^{i\pi B \text{wt}(\Psi_\alpha)} \rho^{\text{wt}(\Psi_\alpha)} \Psi_{-\alpha}(\mathbf{l}, \mathbf{k}). \end{aligned} \quad (50)$$

4 Torus Intertwining n -Point Functions

4.1 Torus Intertwining n -Point Functions for \mathcal{M}

In this section we compute the torus intertwining n -point functions for \mathcal{M} . In the next section we will use these functions to define and compute the partition and n -point functions on a genus two Riemann surface in the ρ formalism. Define the ‘square-bracket’ vertex operator [Z1]

$$\mathcal{Y}[u \otimes e^\alpha, z] = \mathcal{Y}(q_z^{L(0)}(u \otimes e^\alpha), q_z - 1),$$

for $u \otimes e^\alpha \in \mathcal{M}$ and where $q_z = e^z$. As in [Z1] we find that $(V, \mathcal{Y}[\cdot, \cdot], \mathbf{1}, \tilde{\omega})$ and $(V, \mathcal{Y}(\cdot, \cdot), \mathbf{1}, \omega)$ are isomorphic generalized VOAs with $\tilde{\omega} = \omega - \frac{1}{24}\mathbf{1}$, a new conformal vector with vertex operator

$$Y[\tilde{\omega}, z] = \sum_{n \in \mathbb{Z}} L[n] z^{-n-2}.$$

We let $\text{wt}[u \otimes e^\alpha] = \text{wt}[u] + \frac{1}{2}\alpha^2$ denote the weight of an $L[0]$ homogeneous vector $u \otimes e^\alpha$.

Let $M \otimes e^\alpha$ be an irreducible M -module for some $\alpha \in \mathbb{C}$ with torus partition function

$$Z_\alpha^{(1)}(q) = \text{Tr}_{M \otimes e^\alpha} q^{L(0)-1/24} = \frac{q^{\frac{1}{2}\alpha^2}}{\eta(q)},$$

where $\eta(q) = q^{1/24} \prod_{n \geq 1} (1 - q^n)$ is the Dedekind eta for modular parameter q . We also define the 1-point correlation function for $u \in M$ on $M \otimes e^\alpha$ by

$$Z_\alpha^{(1)}(u; q) = \text{Tr}_{M \otimes e^\alpha} (o(u) q^{L(0)-1/24}),$$

where $o(u) = u(1 - \text{wt}(u))$ is the Virasoro level preserving ‘‘zero mode’’ for u of weight $\text{wt}(u)$. In general we define the genus one intertwining n -point correlation function on $M \otimes e^\alpha$ for n vectors $u_1 \otimes e^{\beta_1}, \dots, u_n \otimes e^{\beta_n} \in \mathcal{M}$ by

$$\begin{aligned} & Z_\alpha^{(1)}(u_1 \otimes e^{\beta_1}, z_1; \dots; u_n \otimes e^{\beta_n}, z_n; q) \\ &= \text{Tr}_{M \otimes e^\alpha} \left(\mathcal{Y} \left(q_1^{L(0)}(u_1 \otimes e^{\beta_1}), q_1 \right) \dots \mathcal{Y} \left(q_n^{L(0)}(u_n \otimes e^{\beta_n}), q_n \right) q^{L(0)-1/24} \right), \end{aligned} \quad (51)$$

for formal $q_i = e^{z_i}$ with $i = 1, \dots, n$. Since $e^{\beta \hat{q}} M \otimes e^\alpha = M \otimes e^{\alpha+\beta}$ it follows that the n -point function vanishes when $\sum_{i=1}^n \beta_i \neq 0$.

We next describe a natural generalization of previous results in [MT1] and [MTZ]. Firstly, consider the n -point functions for n highest weight vectors $\mathbf{1} \otimes e^{\beta_i}$, which we abbreviate below to e^{β_i} , for $i = 1, \dots, n$.

Proposition 4.1 *For $\sum_{i=1}^n \beta_i = 0$ then*

$$Z_\alpha^{(1)}(e^{\beta_1}, z_1; \dots; e^{\beta_n}, z_n; q) = \frac{q^{\frac{1}{2}\alpha^2}}{\eta(\tau)} \exp \left(\alpha \sum_{i=1}^n \beta_i z_i \right) \prod_{1 \leq r < s \leq n} K(z_{rs}, \tau)^{\beta_r \beta_s}, \quad (52)$$

where $z_{rs} = z_r - z_s$ and $K(z, \tau)$ is the genus one prime form (11).

The proof appears in Appendix 6.2 and is a natural generalization of results developed in [MT1] and [MTZ].

Remark 4.2 Let $F_\alpha^\beta(\mathbf{z}) = Z_\alpha^{(1)}(e^{\beta_1}, z_1; \dots; e^{\beta_n}, z_n; q)$ of (52).

(i) $F_\alpha^\beta(\mathbf{z})$ is a function of the differences $z_{ij} = z_i - z_j$. Thus we may choose $z_n = 0$ wlog.

(ii) $F_\alpha^\beta(\mathbf{z})$ is analytic on $\{z_i \in \mathbb{C}, \tau \in \mathbb{H} : |z_1| > \dots > |z_n|\}$.

(iii) Note that

$$\begin{aligned} & (z_1 - z_2)^{-\beta_1\beta_2} Z_\alpha^{(1)}(e^{\beta_1}, z_1; e^{\beta_2}, z_2; \dots; e^{\beta_n}, z_n; q) \\ &= (z_2 - z_1)^{-\beta_1\beta_2} Z_\alpha^{(1)}(e^{\beta_2}, z_2; e^{\beta_1}, z_1; \dots; e^{\beta_n}, z_n; q), \end{aligned}$$

in compliance with (38) and (43) (for $N = 0$). Similar identities hold under all label permutations. Thus $F_\alpha^\beta(\mathbf{z})$ can be extended to all $z_i \neq z_j$ for $i \neq j$ up to branch choices.

We may similarly apply the arguments mutatis mutandis of Proposition 1 of [MT1] to obtain a closed form for the general n -point function (51). In particular, due to (41), we may apply standard genus one Zhu recursion theory [Z1] to reduce (51) to an explicit multiple of (52) to find

Proposition 4.3 For $\sum_{i=1}^n \beta_i = 0$ then

$$\begin{aligned} & Z_\alpha^{(1)}(u_1 \otimes e^{\beta_1}, z_1; \dots; u_n \otimes e^{\beta_n}, z_n; q) \\ &= Q_\alpha^{\beta_1, \dots, \beta_n}(u_1, z_1; \dots; u_n, z_n; q) Z_\alpha^{(1)}(e^{\beta_1}, z_1; \dots; e^{\beta_n}, z_n; q), \end{aligned}$$

where $Q_\alpha^{\beta_1, \dots, \beta_n}(u_1, z_1; \dots; u_n, z_n; q)$ is an explicit sum of elliptic and quasi-modular forms (see [MT1] for details).

4.2 Torus Intertwined n -Point Functions for $V_{\mathbb{Z}}$

Let g_1, h_1 be commuting automorphisms of $V_{\mathbb{Z}}$ defined by

$$\sigma f_1 = e^{2\pi i \beta_1 a(0)}, \quad \sigma g_1 = e^{-2\pi i \alpha_1 a(0)}, \quad (53)$$

where $\sigma = e^{\pi i a(0)}$ is the fermion number automorphism. We assume that $\alpha_1, \beta_1 \in \mathbb{R}$ so that

$$\theta_1 = -e^{-2\pi i \beta_1}, \quad \phi_1 = -e^{2\pi i \alpha_1},$$

are of unit modulus (to ensure convergence of Szegő kernels below).

Following Remark 4.2 (i) we consider torus orbifold intertwining $n + 2$ -point functions for the σ_{g_1} -twisted module $V_{\mathbb{Z}+\alpha_1} = \bigoplus_{m \in \mathbb{Z}} M \otimes e^{m+\alpha_1}$ for $V_{\mathbb{Z}}$ defined by:

$$\begin{aligned}
& Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (u_1 \otimes e^{m_1}, z_1; \dots, u_n \otimes e^{m_n}, z_n; v_1 \otimes e^{n_1+\kappa}, w; v_2 \otimes e^{n_2-\kappa}, 0; q) \\
&= \text{Tr}_{V_{\mathbb{Z}+\alpha_1}} \left(\sigma_{f_1} Y \left(q_1^{L(0)}(u_1 \otimes e^{m_1}), q_1 \right) \dots Y \left(q_n^{L(0)}(u_n \otimes e^{m_n}), q_n \right) \right) \cdot \\
&\quad \mathcal{Y} \left(q_w^{L(0)}(v_1 \otimes e^{n_1+\kappa}), q_w \right) \mathcal{Y} \left(v_2 \otimes e^{n_2-\kappa}, 1 \right) q^{L(0)-1/24} \\
&= \sum_{\mu \in \mathbb{Z}+\alpha_1} e^{2\pi i \mu \beta_1} Z_{\mu}^{(1)} (u_1 \otimes e^{m_1}, z_1; \dots; u_n \otimes e^{m_n}, z_n; \\
&\quad v_1 \otimes e^{n_1+\kappa}, w; v_2 \otimes e^{n_2-\kappa}, 0; q), \tag{54}
\end{aligned}$$

where $m_i, n_1, n_2 \in \mathbb{Z}$ for $\kappa \in (-\frac{1}{2}, \frac{1}{2})$ and $q_w = e^w$. The points $w, 0$ will later be identified with the centres of the sewn annuli in the genus two ρ -sewing scheme described in Section 2.2.

In Propositions 14 and 15 of [MTZ] all torus orbifold n -point functions for vectors in $V_{\mathbb{Z}}$ are computed by means of a generating function. In that analysis we made use of an explicit Zhu reduction formula developed for a VOSA with real grading. However, in the present case we do not have an intertwining Zhu reduction formula because of the absence of a generalized VOA commutator formula [DL]. Instead, here we adopt an alternative approach for computing the intertwining n -point functions by exploiting the bosonized formalism. Thus for example, we find

$$\begin{aligned}
Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (e^{\kappa}, w; e^{-\kappa}, 0; q) &= \sum_{\mu \in \mathbb{Z}+\alpha_1} e^{2\pi i \mu \beta_1} Z_{\mu}^{(1)} (e^{\kappa}, w; e^{-\kappa}, 0; q) \\
&= \frac{1}{\eta(q)} \frac{\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (\kappa w, \tau)}{K(w, \tau)^{\kappa^2}}, \tag{55}
\end{aligned}$$

for genus one theta series (1). More generally, we define a generating function

for all $n + 2$ -point functions (54) by the following formal differential form

$$\begin{aligned} & \mathcal{G}_n^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (x_1, y_1, \dots, x_n, y_n) \\ & \equiv Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (\psi^+, x_1; \psi^-, y_1; \dots; \psi^+, x_n; \psi^-, y_n; e^\kappa, w; e^{-\kappa}, 0; q) \prod_{i=1}^n dx_i^{\frac{1}{2}} dy_i^{\frac{1}{2}}, \end{aligned} \quad (56)$$

for $V_{\mathbb{Z}}$ generators $\psi^\pm = e^{\pm 1}$ alternatively inserted at x_i, y_i for $i = 1, \dots, n$.

Proposition 4.4 *The generating form (56) is given by*

$$\mathcal{G}_n^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (x_1, y_1, \dots, x_n, y_n) = Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (e^\kappa, w; e^{-\kappa}, 0; q) \det S_\kappa, \quad (57)$$

where S_κ denotes the $n \times n$ matrix with components $S_\kappa(x_i, y_j)$ for $i, j = 1, \dots, n$ for Szegő kernel (14).

Proof. Proposition 4.1 implies that

$$\begin{aligned} & \mathcal{G}_n^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (x_1, y_1, \dots, x_n, y_n) \\ & = \frac{1}{\eta(\tau) K(w, \tau)^{\kappa^2}} \vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} \left(\sum_{m=1}^n (x_m - y_m) + \kappa w, \tau \right) \\ & \quad \cdot \frac{\prod_{1 \leq i < j \leq n} K(x_i - x_j, \tau) K(y_i - y_j, \tau)}{\prod_{1 \leq i, j \leq n} K(x_i - y_j, \tau)} \prod_{1 \leq i, j \leq n} \left[\frac{K(x_i - w, \tau) K(y_j, \tau)}{K(x_i, \tau) K(y_j - w, \tau)} \right]^\kappa dx_i^{\frac{1}{2}} dy_j^{\frac{1}{2}}. \end{aligned}$$

Next we recall the Frobenius identity (e.g. [F], [MTZ])

$$\frac{\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} \left(\sum_{m=1}^n (x_m - y_m), \tau \right) \prod_{1 \leq i < j \leq n} K(x_i - x_j, \tau) K(y_i - y_j, \tau)}{\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (0, \tau) \prod_{1 \leq i, j \leq n} K(x_i - y_j, \tau)} = \det P_1,$$

where P_1 denotes the $n \times n$ matrix with twisted Weierstrass function components $P_1 \begin{bmatrix} \alpha_1 \\ \phi_1 \end{bmatrix} (x_i, y_j)$ of (13) for $i, j = 1, \dots, n$. Noting that $\vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (x + \kappa w, \tau) =$

$\vartheta \left[\begin{smallmatrix} \alpha_1 \\ \beta_1 - \frac{\kappa w}{2\pi i} \end{smallmatrix} \right] (x, \tau)$ we obtain

$$\begin{aligned} & \vartheta \left[\begin{smallmatrix} \alpha_1 \\ \beta_1 \end{smallmatrix} \right] \left(\sum_{m=1}^n (x_m - y_m) + \kappa w, \tau \right) \frac{\prod_{1 \leq i < j \leq n} K(x_i - x_j, \tau) K(y_i - y_j, \tau)}{\prod_{1 \leq i, j \leq n} K(x_i - y_j, \tau)} \\ &= \vartheta \left[\begin{smallmatrix} \alpha_1 \\ \beta_1 \end{smallmatrix} \right] (\kappa w, \tau) \det \left[\begin{array}{cc} \vartheta \left[\begin{smallmatrix} \alpha_1 \\ \beta_1 \end{smallmatrix} \right] (x_i - y_j + \kappa w, \tau) & 1 \\ \vartheta \left[\begin{smallmatrix} \alpha_1 \\ \beta_1 \end{smallmatrix} \right] (\kappa w, \tau) & K(x_i - y_j, \tau) \end{array} \right]. \end{aligned}$$

The result follows on absorbing the $\left[\frac{K(x_i - w, \tau) K(y_j, \tau)}{K(x_i, \tau) K(y_j - w, \tau)} \right]^\kappa dx_i^{\frac{1}{2}} dy_j^{\frac{1}{2}}$ factors into the determinant and using (55). \square

Finally, we obtain the following generalization of Proposition 15 of [MTZ] concerning the generating properties of (56).

Proposition 4.5 $\mathcal{G}_m^{(1)} \left[\begin{smallmatrix} f_1 \\ g_1 \end{smallmatrix} \right] (x_1, y_1, \dots, x_m, y_m)$ is a generating function for all torus orbifold intertwining $n + 2$ -point functions (54). In particular, for a pair of square bracket mode twisted Fock vectors (34)

$$\begin{aligned} \Psi_\kappa[\mathbf{k}_1, \mathbf{l}_2] &= e^{\kappa \hat{q}} \psi^+[-k_{11}] \dots \psi^+[-k_{1s_1}] \psi^-[-l_{21}] \dots \psi^-[-l_{2t_2}] \mathbf{1} \\ \Psi_{-\kappa}[\mathbf{k}_2, \mathbf{l}_1] &= e^{-\kappa \hat{q}} \psi^+[-k_{21}] \dots \psi^+[-k_{2s_2}] \psi^-[-l_{11}] \dots \psi^-[-l_{1t_1}] \mathbf{1}, \end{aligned}$$

for $p = s_1 + s_2 = t_1 + t_2 > 0$ then

$$\begin{aligned} & Z_{V_Z}^{(1)} \left[\begin{smallmatrix} f_1 \\ g_1 \end{smallmatrix} \right] (\Psi_\kappa[\mathbf{k}_1, \mathbf{l}_2], w; \Psi_{-\kappa}[\mathbf{k}_2, \mathbf{l}_1], 0; q) \\ &= \epsilon Z_{V_Z}^{(1)} \left[\begin{smallmatrix} f_1 \\ g_1 \end{smallmatrix} \right] (e^\kappa, w; e^{-\kappa}, 0; q) \det C_{ab}(\mathbf{k}_a, \mathbf{l}_b) \end{aligned} \quad (58)$$

where

$$\epsilon = (-1)^{(t_1 + s_2)t_2 + \lfloor \frac{1}{2}p \rfloor} e^{i\pi B\kappa(s_2 - t_1)},$$

for the odd integer B fixed in (43) and

$$C_{ab}(\mathbf{k}_a, \mathbf{l}_b) = \begin{bmatrix} C_{11}(\mathbf{k}_1, \mathbf{l}_1) & C_{12}(\mathbf{k}_1, \mathbf{l}_2) \\ C_{21}(\mathbf{k}_2, \mathbf{l}_1) & C_{22}(\mathbf{k}_2, \mathbf{l}_2) \end{bmatrix}$$

is a $p \times p$ block matrix with components $C_{ab}(k_{ai_a}, l_{bj_b})$ for $i_a = 1, \dots, s_a$ and $j_b = 1, \dots, t_b$ for $a, b = 1, 2$ for S_κ expansion coefficients (16).

Proof. Consider the generating function for ψ^+ and ψ^- alternatively inserted at $x_{21} + w, \dots, x_{2s_2} + w, x_{11}, \dots, x_{1s_1}$ and $y_{21} + w, \dots, y_{2t_2} + w, y_{11}, \dots, y_{2t_1}$ respectively. Reordering the vertex operators and using generalized locality (43) we obtain

$$\begin{aligned} \widehat{\epsilon} Z_{V_{\mathbb{Z}}}^{(1)} \left[\begin{matrix} f_1 \\ g_1 \end{matrix} \right] & (\psi^+, x_{21} + w; \dots; \psi^+, x_{2s_2} + w; \psi^-, y_{21} + w; \dots; \psi^-, y_{2t_2} + w; e^\kappa, w; \\ & \psi^+, x_{11}; \dots; \psi^+, x_{1s_1}; \psi^-, y_{11}; \dots; \psi^-, y_{1t_1}; e^{-\kappa}, 0; q) \prod_{i_a, j_b} dx_{ai_a}^{\frac{1}{2}} dy_{bj_b}^{\frac{1}{2}}, \end{aligned} \quad (59)$$

for $\widehat{\epsilon} = (-1)^{s_2 t_2 + \lfloor \frac{1}{2} p \rfloor} e^{-i\pi B \kappa (s_2 - t_1)}$. Generalized associativity (40) implies that

$$\begin{aligned} & \prod_{i,j}^{s_1} (x_{2i} + w)^{M_i - \kappa} (y_{2j} + w)^{N_j + \kappa} \mathcal{Y}[\psi^+, x_{21} + w] \dots \mathcal{Y}[\psi^-, y_{2t_2} + w] \mathcal{Y}[e^\kappa, w] \\ &= \prod_{i,j}^{s_1} (w + x_{2i})^{M_i - \kappa} (w + y_{2j})^{N_j + \kappa} \mathcal{Y}[\mathcal{Y}[\psi^+, x_{21}] \dots \mathcal{Y}[\psi^-, y_{2t_2}] e^\kappa, w], \end{aligned}$$

for $M_i, N_j \gg 0$. But using (90) and $\Delta(\kappa, z)\psi^\pm = z^{\pm\kappa}\psi^\pm$ we find that

$$\begin{aligned} & Y[\psi^+, x_{21}] \dots Y[\psi^+, x_{2s_1}] Y[\psi^-, y_{21}] \dots Y[\psi^-, y_{2t_2}] e^\kappa \\ &= \prod_{i=1}^{s_1} x_{2i}^\kappa \prod_{j=1}^{t_2} y_{2j}^{-\kappa} \cdot e^{\kappa \widehat{[q]}} Y[\psi^+, x_{21}] \dots Y[\psi^+, x_{2s_1}] Y[\psi^-, y_{21}] \dots Y[\psi^-, y_{2t_2}] \mathbf{1} \\ &= \sum_{\mathbf{k}_1 \in \mathbb{Z}^{s_1}, \mathbf{l}_2 \in \mathbb{Z}^{t_2}} \prod_{i=1}^{s_1} x_{2i}^{k_{1i} + \kappa - 1} \prod_{j=1}^{t_2} y_{2j}^{l_{2j} - \kappa - 1} \Psi_\kappa[\mathbf{k}_1, \mathbf{l}_2], \end{aligned}$$

generates the Fock vector $\Psi_\kappa[\mathbf{k}_1, \mathbf{l}_2]$ as the coefficient of $\prod_{i=1}^{s_1} x_{2i}^{k_{1i} + \kappa - 1} \prod_{j=1}^{t_2} y_{2j}^{l_{2j} - \kappa - 1}$ (and where $[a[n], \widehat{[q]}] = \delta_{n0}$ is the square bracket version of (32)). Similarly $\Psi_{-\kappa}[\mathbf{k}_2, \mathbf{l}_1]$ is the coefficient of $\prod_{i=1}^{s_2} x_{1i}^{k_{2i} - \kappa - 1} \prod_{j=1}^{t_1} y_{1j}^{l_{1j} + \kappa - 1}$ in

$$Y[\psi^+, x_{11}] \dots Y[\psi^+, x_{1s_2}] Y[\psi^-, y_{11}] \dots Y[\psi^-, y_{1t_1}] e^{-\kappa}.$$

(58) follows by reading off the coefficient in the corresponding expansion (16) of the Szegő kernels in the determinant of (57). Finally, interchanging the first t_1 and second t_2 columns of the resulting determinant gives

$(-1)^{t_1 t_2} \det C_{ab}(\mathbf{k}_a, \mathbf{l}_b)$. Thus $\epsilon = (-1)^{t_1 t_2} / \widehat{\epsilon}$ as given. In general, we can expand (57) in appropriate parameters much as in Proposition 15 of [MTZ] to obtain any $n+2$ -point function for $V_{\mathbb{Z}}$ -Fock vectors inserted at z_1, \dots, z_n for $z_i \neq 0, w$ together with the above pair of twisted Fock vectors. \square

5 Genus Two Twisted Partition and n -Point Functions

We now arrive at the main results of this paper, namely, the computation of the genus two twisted partition and n -point function for $V_{\mathbb{Z}}$ on the genus two Riemann surface formed by self-sewing a torus as described in Section 2.2. Our definitions are a natural extension of those for a VOA introduced in [MT3]. In the present case, we consider genus two twisted partition and n -point function which depend on four $V_{\mathbb{Z}}$ -automorphisms:

$$\begin{aligned} \sigma f_1 &= e^{2\pi i \beta_1 a(0)}, & \sigma g_1 &= e^{-2\pi i \alpha_1 a(0)}, \\ \sigma f_2 &= e^{2\pi i \beta_2 a(0)}, & \sigma g_2 &= e^{-2\pi i \kappa a(0)}, \end{aligned} \quad (60)$$

for real $\alpha_1, \beta_1, \beta_2$ and $\kappa \in (-\frac{1}{2}, \frac{1}{2})$. We let $f = (f_1, f_2)$ and $g = (g_1, g_2)$ denote pairs of automorphisms. We also define unit modulus parameters

$$\begin{aligned} \theta_1 &= -e^{-2\pi i \beta_1}, & \phi_1 &= -e^{2\pi i \alpha_1}, \\ \theta_2 &= -e^{-2\pi i \beta_2}, & \phi_2 &= -e^{2\pi i \kappa}, \end{aligned} \quad (61)$$

later identified with the multipliers (3) of a genus two Szegő kernel below.

Following [MT3] we define the continuous orbifold genus two n -point function for $u_i \otimes e^{m_i} \in V_{\mathbb{Z}}$ inserted at z_i on the sewn genus two Riemann surface in the ρ sewing scheme as the following infinite sum of genus one intertwined $n+2$ point functions:

$$\begin{aligned} & Z_{V_{\mathbb{Z}}}^{(2)} \left[\begin{matrix} f \\ g \end{matrix} \right] (u_1 \otimes e^{m_1}, z_1; \dots, u_n \otimes e^{m_n}, z_n; \tau, w, \rho) \\ &= \sum_{\Psi_{\kappa}} Z_{V_{\mathbb{Z}}}^{(1)} \left[\begin{matrix} f_1 \\ g_1 \end{matrix} \right] (u_1 \otimes e^{m_1}, z_1; \dots, u_n \otimes e^{m_n}, z_n; \sigma f_2 \Psi_{\kappa}, w; \overline{\Psi}_{\kappa}, 0; q). \end{aligned} \quad (62)$$

The sum is taken over the square bracket Fock basis $\Psi_{\kappa} \in V_{\mathbb{Z}+\kappa}$ with square bracket λ -dual $\overline{\Psi}_{\kappa}$ for $\lambda = (e^{i\pi B} \rho)^{\frac{1}{2}}$ of (46) and with σf_2 acting on Ψ_{κ} .

5.1 Genus Two Twisted Partition Function

The partition or 0–point function is expressible in terms of the basic genus one twisted 2–point function (55) and the infinite Szegő moment matrix $T = T(\tau, w, \rho | \theta_1, \theta_2, \phi_1, \kappa)$ of (23) as follows:

Theorem 5.1 *The genus two continuous orbifold partition function for $V_{\mathbb{Z}}$ on a Riemann surface in the ρ -sewing formalism is convergent on \mathcal{D}^p and is given by*

$$Z_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\tau, w, \rho) = e^{2i\pi\beta_2\kappa} (e^{i\pi B} \rho)^{\frac{1}{2}\kappa^2} Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (e^\kappa, w; e^{-\kappa}, 0; q) \cdot \det(I - T). \quad (63)$$

In order to prove Theorem 5.1 we first recall Proposition 2 of [TZ2]. Let R be an $P \times P$ matrix and let $\mathbf{m} = (m_1, \dots, m_p)$ denote p ordered subindices with $1 \leq m_1 < \dots < m_p \leq P$. We refer to \mathbf{m} as an P -subindex of length p . Let

$$R(\mathbf{m}, \mathbf{m}) = (R(m_r, m_s)),$$

for $r, s = 1, \dots, p$ denote the $p \times p$ submatrix of R . We define $R(\mathbf{m}, \mathbf{m}) = 1$ in the degenerate case $p = 0$ and let I denote the identity matrix. We have

Proposition 5.2

$$\det(I + R) = \sum_{p=0}^P \sum_{\mathbf{m}} \det R(\mathbf{m}, \mathbf{m}), \quad (64)$$

where the inner sum runs over all P -subindices of length p . \square

Proof of Theorem 5.1. Recall the Fock vector $\Psi_\kappa = e^{\kappa\hat{q}}\Psi$ with

$$\Psi = \Psi[\mathbf{k}, \mathbf{l}] = \psi^+[-k_1] \dots \psi^+[-k_s] \psi^-[-l_1] \dots \psi^-[-l_t] \mathbf{1},$$

has Li–Z dual $\bar{\Psi}_\kappa[\mathbf{k}, \mathbf{l}] = \epsilon_1 \rho^{\text{wt}[\Psi_\kappa]} \Psi_{-\kappa}[\mathbf{l}, \mathbf{k}]$ where

$$\epsilon_1 = (-1)^{st + \lfloor \text{wt}[\Psi] \rfloor} e^{i\pi B \text{wt}[\Psi_\kappa]}, \quad (65)$$

for $\text{wt}[\Psi_\kappa] = \text{wt}[\Psi] + \kappa(s - t) + \frac{1}{2}\kappa^2$. Furthermore,

$$\sigma f_2 \Psi_\kappa = (-\theta_2)^{t-s} e^{2i\pi\beta_2\kappa} \Psi_\kappa,$$

from (60) and (61). Thus we apply (58) of Proposition (4.5) with $\mathbf{k} = \mathbf{k}_1 = \mathbf{l}_1$ for $s = s_1 = t_1$ and $\mathbf{l} = \mathbf{k}_2 = \mathbf{l}_2$ for $t = s_2 = t_1$ and $p = s + t$ to obtain

$$\begin{aligned} Z_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\tau, w, \rho) &= e^{2i\pi\beta_2\kappa} Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (e^\kappa, w; e^{-\kappa}, 0; q) \\ &\cdot \sum_{\mathbf{k}, \mathbf{l}} \epsilon_1 \epsilon_2 (-\theta_2)^{t-s} \rho^{\text{wt}[\Psi_\kappa]} \det \begin{bmatrix} C_{11}(\mathbf{k}, \mathbf{k}) & C_{12}(\mathbf{k}, \mathbf{l}) \\ C_{21}(\mathbf{l}, \mathbf{k}) & C_{22}(\mathbf{l}, \mathbf{l}) \end{bmatrix}, \end{aligned}$$

where

$$\epsilon_2 = (-1)^{t+st+\lfloor \frac{1}{2}p \rfloor} e^{i\pi B\kappa(t-s)}. \quad (66)$$

Noting that $(-1)^{\lfloor \text{wt}[\Psi] \rfloor} e^{i\pi B \text{wt}[\Psi]} = (-1)^{\lfloor \frac{1}{2}p \rfloor} e^{\frac{1}{2}i\pi Bp}$ we find

$$\epsilon_1 \epsilon_2 (-\theta_2)^{t-s} \rho^{\text{wt}[\Psi_\kappa]} = (-\xi\theta_2^{-1})^s (\xi\theta_2)^t \rho^{\text{wt}[\Psi]+\kappa(s-t)} (e^{i\pi B} \rho)^{\frac{1}{2}\kappa^2},$$

for $\xi = e^{\frac{1}{2}i\pi B}$ of (22). We may absorb all but the last of these factors into the $p \times p$ determinant to find

$$\begin{aligned} &(-\xi\theta_2^{-1})^s (\xi\theta_2)^t \rho^{\text{wt}[\Psi]+\kappa(s-t)} \det \begin{bmatrix} C_{11}(k_i, k_j) & C_{12}(k_i, l_{j'}) \\ C_{21}(l_{i'}, k_j) & C_{22}(l_{i'}, l_{j'}) \end{bmatrix} \\ &= \det \begin{bmatrix} -\xi\theta_2^{-1} \rho^{\frac{1}{2}(k_i+k_j+2\kappa-1)} C_{11}(k_i, k_j) & \xi\theta_2 \rho^{\frac{1}{2}(k_i+l_{j'}-1)} C_{12}(k_i, l_{j'}) \\ -\xi\theta_2^{-1} \rho^{\frac{1}{2}(l_{i'}+k_j-1)} C_{21}(l_{i'}, k_j) & \xi\theta_2 \rho^{\frac{1}{2}(l_{i'}+l_{j'}-2\kappa-1)} C_{22}(l_{i'}, l_{j'}) \end{bmatrix} \\ &= \det(-T(\mathbf{m}, \mathbf{m})), \end{aligned}$$

for T of (23) with $i, j = 1, \dots, s$ and $i', j' = 1, \dots, t$ and where $\mathbf{m} = (\mathbf{k}, \mathbf{l})$ i.e. $m_i = k_i$ and $m_{i'+s} = l_{i'}$. But \mathbf{m} is a P -subindex of length p for some sufficiently large P so that we may apply Proposition 5.2 for sufficiently large P by truncating the partition function to any finite order in ρ to obtain the result. Finally, convergence on \mathcal{D}^ρ is guaranteed by Theorem 2.1. \square

5.2 The Genus Two n -Point Generating Form

Similarly to the genus one situation (56), we define a genus two continuous orbifold generating differential form by

$$\begin{aligned} &\mathcal{G}_n^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (x_1, y_1, \dots, x_n, y_n) \\ &= Z_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\psi^+, x_1; \psi^-, y_1; \dots, \psi^+, x_n; \psi^-, y_n; \tau, w, \rho) \prod_{i=1}^n dx_i^{\frac{1}{2}} dy_i^{\frac{1}{2}}. \end{aligned} \quad (67)$$

Recalling (14), (20), (21) and (25) we define matrices

$$\begin{aligned} S^{(2)} &= (S^{(2)}(x_i, y_j)), \quad S_\kappa = (S_\kappa(x_i, y_j)), \\ H &= ((h(x_i))(k, a)), \quad \overline{H}^T = ((\overline{h}(y_i))(l, b))^T. \end{aligned} \quad (68)$$

$S^{(2)}$ and S_κ are finite matrices indexed by $i, j = 1, \dots, n$; H is semi-infinite with n rows indexed by i and columns indexed by $k \geq 1$ and $a = 1, 2$ and \overline{H}^T is semi-infinite with rows indexed by $l \geq 1$ and $b = 1, 2$ and with n columns indexed by j .

We can now state the second main result of this paper.

Theorem 5.3 *The generating form (67) is given by*

$$\mathcal{G}_n^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (x_1, y_1, \dots, x_n, y_n) = Z_{V_z}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\tau, w, \rho) \det S^{(2)}, \quad (69)$$

for $n \times n$ matrix $S^{(2)}$ of (68) and $Z_{V_z}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\tau, w, \rho)$ is the genus two twisted partition function (63).

Normalized relative to the genus two partition function, the 2-point function for ψ^+ and ψ^- is thus given by the Szegő kernel and more generally, the generating function is a Szegő kernel determinant. This agrees with the *assumed* form in [R] or as found by string theory methods using a Schottky parametrization in [DVPFHLS] and in the genus two ϵ -sewing scheme [TZ2].

In order to prove Theorem 5.3 we firstly note that similarly to Proposition 3 in [TZ2] we have

Proposition 5.4

$$\det \begin{bmatrix} S_\kappa & -\xi H D^{\theta_2} \\ \overline{H}^T & I - T \end{bmatrix} = \det S^{(2)} \det(I - T), \quad (70)$$

with T, D^{θ_2} of (23) and (24). \square

We also note the following generalization of Proposition 5.2. Let S, U, V, R be $n \times n, n \times P, P \times n$ and $P \times P$ matrices respectively. For \mathbf{m} , a P -subindex of length p , we let

$$\begin{aligned} U(\mathbf{m}) &= (U(i, m_s)), \quad i = 1, \dots, n; \quad s = 1, \dots, p, \\ V(\mathbf{m}) &= (V(m_r, j)), \quad r = 1, \dots, p; \quad j = 1, \dots, n. \end{aligned}$$

We then find

Proposition 5.5

$$\det \begin{bmatrix} S & U \\ V & I + R \end{bmatrix} = \sum_{p=0}^P \sum_{\mathbf{m}} \det \begin{bmatrix} S & U(\mathbf{m}) \\ V(\mathbf{m}) & R(\mathbf{m}, \mathbf{m}) \end{bmatrix}, \quad (71)$$

where the inner sum runs over all P -subindices of length p . \square

Proof of Theorem 5.3. Repeating the notation of the proof of Theorem 5.1

we find that $G_n^{(2)}(x_1, \dots, y_n) \prod_{i=1}^n dx_i^{-\frac{1}{2}} dy_i^{-\frac{1}{2}}$ is

$$e^{2i\pi\beta_2\kappa} \sum_{\mathbf{k}, \mathbf{l}} \epsilon_1(-\theta_2)^{t-s} \rho^{\text{wt}[\Psi_\kappa]} Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (\psi^+, x_1; \dots; \psi^-, y_n; \Psi_\kappa[\mathbf{k}, \mathbf{l}], w; \Psi_{-\kappa}[\mathbf{l}, \mathbf{k}], 0; q),$$

with ϵ_1 of (65). The expansion technique described in Proposition 4.5 implies

$$\begin{aligned} & Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (\psi^+, x_1; \dots; \psi^-, y_n; \Psi_\kappa[\mathbf{k}, \mathbf{l}], w; \Psi_{-\kappa}[\mathbf{l}, \mathbf{k}], 0; q) \prod_{i=1}^n dx_i^{\frac{1}{2}} dy_i^{\frac{1}{2}} \\ &= \epsilon_2 Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (e^\kappa, w; e^{-\kappa}, 0; q) \det \begin{bmatrix} S_\kappa(x_i, y_j) & d_1(x_i, \mathbf{k}) & d_2(x_i, \mathbf{l}) \\ \bar{d}_1(y_j, \mathbf{k}) & C_{11}(\mathbf{k}, \mathbf{k}) & C_{12}(\mathbf{k}, \mathbf{l}) \\ \bar{d}_2(y_j, \mathbf{l}) & C_{21}(\mathbf{l}, \mathbf{k}) & C_{22}(\mathbf{l}, \mathbf{l}) \end{bmatrix}, \end{aligned}$$

for ϵ_2 of (66) and $d_a(x, k), \bar{d}_a(y, k)$ of (17) and (18). Absorbing various factors as before we find

$$\begin{aligned} & (-\xi)^p (-1)^t \theta_2^{t-s} \rho^{\text{wt}[\Psi] + \kappa(s-t)} \det \begin{bmatrix} S_\kappa(x_i, y_j) & d_1(x_i, \mathbf{k}) & d_2(x_i, \mathbf{l}) \\ \bar{d}_1(y_j, \mathbf{k}) & C_{11}(\mathbf{k}, \mathbf{k}) & C_{12}(\mathbf{k}, \mathbf{l}) \\ \bar{d}_2(y_j, \mathbf{l}) & C_{21}(\mathbf{l}, \mathbf{k}) & C_{22}(\mathbf{l}, \mathbf{l}) \end{bmatrix} \\ &= \det \begin{bmatrix} S_\kappa & -\xi (HD^{\theta_2})(\mathbf{m}) \\ \left(\overline{H}^T\right)(\mathbf{m}) & -T(\mathbf{m}, \mathbf{m}) \end{bmatrix}, \end{aligned}$$

for $\mathbf{m} = (\mathbf{k}, \mathbf{l})$ as before. The result follows on applying Proposition 5.5 and followed by Proposition 5.4. \square

5.3 Bosonization

We may compute the genus two twisted partition and generating functions in an alternative way by use of a bosonic basis $u \otimes e^\nu \in V_{\mathbb{Z}+\kappa}$ for $u \in M$ and

$\nu \in \mathbb{Z} + \kappa$. In particular, we can immediately exploit and extend the results for lattice VOAs in [MT3] in the light of Section 4 above.

Let us firstly recall from [MT2, MT3] the following definitions:

$$\begin{aligned} P_2(\tau, z) &= \wp(\tau, z) + E_2(\tau) \\ &= \frac{1}{z^2} + \sum_{k=2}^{\infty} (k-1)E_k(\tau)z^{k-2}, \end{aligned} \quad (72)$$

for Weierstrass function $\wp(\tau, z)$ and Eisenstein series $E_k(\tau)$. We define $P_{k+1}(\tau, z) = -\frac{1}{k}\partial_z P_k(\tau, z)$ for $k \geq 2$ and for $k, l \geq 1$

$$\begin{aligned} C(k, l, \tau) &= (-1)^{k+1} \frac{(k+l-1)!}{(k-1)!(l-1)!} E_{k+l}(\tau), \\ D(k, l, \tau, z) &= (-1)^{k+1} \frac{(k+l-1)!}{(k-1)!(l-1)!} P_{k+l}(\tau, z), \\ R_{ab}(k, l) &= -\frac{\rho^{(k+l)/2}}{\sqrt{kl}} \begin{bmatrix} D(k, l, \tau, w) & C(k, l, \tau) \\ C(k, l, \tau) & D(l, k, \tau, w) \end{bmatrix}. \end{aligned}$$

Theorems 5.1 and 5.6 of [MT3] tell us that:

Theorem 5.6 *The genus two partition function for the Heisenberg VOA M in the ρ sewing scheme is*

$$Z_M^{(2)}(\tau, w, \rho) = Z_M^{(1)}(\tau) \det(1 - R)^{-1/2}, \quad (73)$$

where $Z_M^{(1)}(\tau) = 1/\eta(q)$ is the genus one partition function. $Z_M^{(2)}(\tau, w, \rho)$ is holomorphic on \mathcal{D}^ρ .

Theorem 6.1 of [MT3] concerns the genus two partition function for a lattice VOA. We obtain a natural generalization for a twisted genus two partition function defined by

$$Z_{\mu, \nu}^{(2)}(\tau, w, \rho) = \sum_{\Psi_\nu \in M \otimes e^\nu} Z_\mu^{(1)}(\Psi_\nu, w; \bar{\Psi}_\nu, 0; q), \quad (74)$$

for $\mu, \nu \in \mathbb{C}$ where the sum is taken over a $M \otimes e^\nu$ basis, $\bar{\Psi}_\nu$ is the square bracket λ -dual and the summand is a genus one 2-point function (51) for \mathcal{M} . We find that

Theorem 5.7

$$Z_{\mu,\nu}^{(2)}(\tau, w, \rho) = e^{i\pi(\mu^2\Omega_{11}+2\mu\nu\Omega_{12}+\nu^2\Omega_{22})} Z_M^{(2)}(\tau, w, \rho), \quad (75)$$

where Ω is the genus two period matrix.

Proof. The proof follows precisely that for Theorem 6.1 of [MT3] by use of a bosonic Fock basis and by applying Propositions 4.1 and 4.3. \square

We are now able to compute the genus two twisted partition function (63) by use of a bosonic basis to obtain

Theorem 5.8 *The genus two twisted partition function for $V_{\mathbb{Z}}$ on a Riemann surface in the ρ -sewing formalism is given by*

$$Z_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\tau, w, \rho) = \vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} (\Omega) Z_M^{(2)}(\tau, w, \rho), \quad (76)$$

for genus two Riemann theta function (1) with characteristics $\alpha_1, \alpha_2 = \kappa, \beta_1, \beta_2$ of (61) and where $Z_M^{(2)}(\tau, w, \rho)$ is the genus two Heisenberg partition function.

Proof. Let $\Psi_\nu = u \otimes e^\nu \in V_{\mathbb{Z}+\kappa}$ with square bracket λ -dual $\bar{\Psi}_\nu$. Then

$$\sigma f_2 \Psi_\nu = e^{2i\pi\nu\beta_2} \Psi_\nu,$$

from (60). Furthermore,

$$Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (\Psi_\nu, w, \bar{\Psi}_\nu, 0; q) = \sum_{\mu \in \mathbb{Z} + \alpha_1} e^{2\pi i \mu \beta_1} Z_\mu^{(1)}(\Psi_\nu, w, \bar{\Psi}_\nu, 0; q).$$

Thus altogether we find

$$\begin{aligned} Z_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\tau, w, \rho) &= \sum_{\mu \in \mathbb{Z} + \alpha_1} \sum_{\nu \in \mathbb{Z} + \kappa} e^{2\pi i \mu \beta_1} e^{2i\pi\nu\beta_2} \sum_{\Psi_\nu \in M \otimes e^\nu} Z_\mu^{(1)}(\Psi_\nu, w, \bar{\Psi}_\nu, 0; q) \\ &= \sum_{\mu \in \mathbb{Z} + \alpha_1} \sum_{\nu \in \mathbb{Z} + \kappa} e^{i\pi(\mu^2\Omega_{11}+2\mu\nu\Omega_{12}+\nu^2\Omega_{22})} e^{2\pi i(\mu\beta_1+\nu\beta_2)} Z_M^{(2)}(\tau, w, \rho) \\ &= \vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} (\Omega) Z_M^{(2)}(\tau, w, \rho). \quad \square \end{aligned}$$

Comparing with the fermionic expression (63) and noting that

$$(e^{i\pi B} \rho)^{\frac{1}{2}\kappa^2} Z_{V_{\mathbb{Z}}}^{(1)} \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} (e^\kappa, w; e^{-\kappa}, 0; q) = \left(\frac{e^{i\pi B} \rho}{K(w, \tau)^2} \right)^{\frac{1}{2}\kappa^2} \frac{1}{\eta(\tau)} \vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (\kappa w, \tau), \quad (77)$$

we obtain a genus two analogue of the classical Jacobi triple product identity for the elliptic theta function (separate to a similar identity shown in [TZ2]) as follows:

Theorem 5.9 *The genus two Riemann theta function on \mathcal{D}^ρ is given by*

$$\vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} (\Omega) = e^{2i\pi\beta_2\kappa} \left(\frac{e^{i\pi B\rho}}{K(w, \tau)^2} \right)^{\frac{1}{2}\kappa^2} \vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (\kappa w, \tau) \det(I - T) \det(I - R)^{\frac{1}{2}}.$$

Since $R, T \rightarrow 0$ as $\rho \rightarrow 0$ then $\vartheta \begin{bmatrix} \alpha \\ \beta \end{bmatrix} (\Omega) \approx e^{2i\pi\beta_2\kappa} \left(\frac{e^{i\pi B\rho}}{K(w, \tau)^2} \right)^{\frac{1}{2}\kappa^2} \vartheta \begin{bmatrix} \alpha_1 \\ \beta_1 \end{bmatrix} (\kappa w, \tau)$ to leading order in ρ as discussed in greater detail in [TZ1]. As suggested in [R], we also remark that Fay's Trisecant Identity at genus two follows by comparing the generating function $\mathcal{G}_n^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (x_1, y_1, \dots, x_n, y_n)$ of (67) to its form in terms of a bosonic basis. We do not give the details here.

5.4 Modular Invariance

We lastly consider the modular properties of the genus two partition (63) and n -point generating (69) functions for the rank two fermion VOSA. In [TZ1] we proved the modular invariance of the genus two Szegő kernel $S^{(2)} \begin{bmatrix} \theta^{(2)} \\ \phi^{(2)} \end{bmatrix} (x, y | \tau, w, \rho)$ of (25) under the action of a particular subgroup $L \subset \text{Sp}(4, \mathbb{Z})$ and verified that (4) holds. We also showed that $\det(I - T)$ is modular invariant under L .

Let us recall the relevant definitions used to describe L [MT2]. Consider the Heisenberg group $\widehat{H} \subset \text{Sp}(4, \mathbb{Z})$ with elements

$$\mu(a, b, c) = \begin{pmatrix} 1 & 0 & 0 & b \\ a & 1 & b & c \\ 0 & 0 & 1 & -a \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (78)$$

\widehat{H} is generated by $A = \mu(1, 0, 0)$, $B = \mu(0, 1, 0)$ and $C = \mu(0, 0, 1)$ with relations $[A, B]C^{-2} = [A, C] = [B, C] = 1$. We also define $\Gamma_1 \subset \text{Sp}(4, \mathbb{Z})$ where $\Gamma_1 \cong \text{SL}(2, \mathbb{Z})$ with elements

$$\gamma_1 = \begin{pmatrix} a_1 & 0 & b_1 & 0 \\ 0 & 1 & 0 & 0 \\ c_1 & 0 & d_1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a_1 d_1 - b_1 c_1 = 1. \quad (79)$$

Together these groups generate $L = \widehat{H} \rtimes \Gamma_1 \subset \mathrm{Sp}(4, \mathbb{Z})$ with center $Z(L) = \langle C \rangle$ where $J = L/Z(L) \cong \mathbb{Z}^2 \rtimes \mathrm{SL}(2, \mathbb{Z})$ is the Jacobi group. From Lemma 15 of [MT2] we find that L acts on $x = (\tau, w, \rho) \in \mathcal{D}^\rho$ of (9) as follows:

$$\mu(a, b, c).x = (\tau, w + 2\pi ia\tau + 2\pi ib, \rho), \quad (80)$$

$$\gamma_1.x = \left(\frac{a_1\tau + b_1}{c_1\tau + d_1}, \frac{w}{c_1\tau + d_1}, \frac{\rho}{(c_1\tau + d_1)^2} \right). \quad (81)$$

Note that the Jacobi group J acts properly on \mathcal{D}^ρ .

The genus two partition function $Z_{V_{\mathbb{Z}}}^{(2)} \left[\begin{smallmatrix} f \\ g \end{smallmatrix} \right] (\tau, w, \rho)$ is not single-valued on \mathcal{D}^ρ due to the $(e^{i\pi B} \rho K(w, \tau)^{-2})^{\frac{1}{2}\kappa^2}$ factor in (63). In Section 6 of [MT2] a detailed description is given of the logarithmic function

$$l(x) = \log \left(-\frac{\rho}{K(\tau, w)^2} \right), \quad x = (\tau, w, \rho) \in \mathcal{D}^\rho. \quad (82)$$

In particular, we describe a covering space $\widehat{\mathcal{D}}^\rho$ with projection $p : \widehat{\mathcal{D}}^\rho \rightarrow \mathcal{D}^\rho$ where $p(\widehat{x}) = (\tau, w, \rho)$ on which l lifts to a single-valued holomorphic function $\widehat{l} : \widehat{\mathcal{D}}^\rho \rightarrow \mathbb{C}$. Furthermore, L acts on $\widehat{\mathcal{D}}^\rho$ with $p(\gamma.\widehat{x}) = \gamma.x$ for all $\gamma \in L$ where [MT2]

$$\widehat{l}(\mu(a, b, c).\widehat{x}) = \widehat{l}(\widehat{x}) + 2\pi ia^2\tau + 2aw + 2\pi i(ab + c), \quad (83)$$

$$\widehat{l}(\gamma_1.\widehat{x}) = \widehat{l}(\widehat{x}) - \frac{1}{2\pi i} \frac{c_1 w^2}{c_1\tau + d_1}. \quad (84)$$

Thus, we may lift $Z_{V_{\mathbb{Z}}}^{(2)} \left[\begin{smallmatrix} f \\ g \end{smallmatrix} \right] (\tau, w, \rho)$ to a holomorphic function $\widehat{Z}_{V_{\mathbb{Z}}}^{(2)} \left[\begin{smallmatrix} f \\ g \end{smallmatrix} \right] (\widehat{x})$ on $\widehat{\mathcal{D}}^\rho$ where the factor $(e^{i\pi B} \rho K(w, \tau)^{-2})^{\frac{1}{2}\kappa^2}$ is lifted to $\exp \left(\frac{1}{2}\kappa^2 \widehat{l}(\widehat{x}) \right)$ and the remaining single-valued terms of $Z_{V_{\mathbb{Z}}}^{(2)} \left[\begin{smallmatrix} f \\ g \end{smallmatrix} \right] (\tau, w, \rho)$ lift trivially.

We next define an action of L on the automorphisms of (60)

$$\mu(a, b, c). \begin{bmatrix} f_1 \\ f_2 \\ g_1 \\ g_2 \end{bmatrix} = \begin{bmatrix} \sigma^b f_1 g_2^b \\ \sigma^{a+b+ab} f_1^a f_2 g_1^b g_2^c \\ \sigma^{-a} g_1 g_2^{-a} \\ g_2 \end{bmatrix}, \quad \gamma_1. \begin{bmatrix} f_1 \\ f_2 \\ g_1 \\ g_2 \end{bmatrix} = \begin{bmatrix} f_1^{a_1} g_1^{b_1} \\ f_2 \\ f_1^{c_1} g_1^{d_1} \\ g_2 \end{bmatrix}. \quad (85)$$

This action is compatible with (6) and it is straightforward to confirm that the defining relations for the generators A, B, C are satisfied. We may also determine the L action on the multipliers θ_i, ϕ_i of (61) merely by replacing

f_i, g_i in (85) by $\theta_i^{-1}, \phi_i^{-1}$, respectively, and σ by -1 .³ We thus define a canonical action of $\gamma \in L$ on the genus two partition function as follows

$$\widehat{Z}_{V_{\mathbb{Z}}}^{(2)} \left[\begin{matrix} f \\ g \end{matrix} \right] \Big|_{\gamma} (\widehat{x}) = \widehat{Z}_{V_{\mathbb{Z}}}^{(2)} \left(\gamma \cdot \left[\begin{matrix} f \\ g \end{matrix} \right] \right) (\gamma \cdot \widehat{x}). \quad (86)$$

In [TZ1] we showed that $\det(I - T)$ is invariant under the modular group L (with L acting on the multipliers θ_i, ϕ_i as in (85)) so we need only prove modularity for the following partition function factor

$$F \left[\begin{matrix} f \\ g \end{matrix} \right] (\widehat{x}) = e^{2i\pi\beta_2\kappa} \exp \left(\frac{1}{2} \kappa^2 \widehat{l}(\widehat{x}) \right) \frac{1}{\eta(\tau)} \vartheta \left[\begin{matrix} \alpha_1 \\ \beta_1 \end{matrix} \right] (\kappa w, \tau).$$

We first analyse the action of the $\Gamma_1 = \text{SL}(2, \mathbb{Z})$ generators $S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and $T = \begin{pmatrix} 1 & \\ 0 & 1 \end{pmatrix}$. We note that

$$\begin{aligned} \vartheta \left[\begin{matrix} -\beta_1 \\ \alpha_1 \end{matrix} \right] \left(\kappa \left(-\frac{w}{\tau} \right), -\frac{1}{\tau} \right) &= (-i\tau)^{\frac{1}{2}} e^{-2\pi i \alpha_1 \beta_1} e^{-i\kappa^2 w^2 / 4\pi\tau} \vartheta \left[\begin{matrix} \alpha_1 \\ \beta_1 \end{matrix} \right] (\kappa w, \tau), \\ \vartheta \left[\begin{matrix} \alpha_1 \\ \beta_1 - \alpha_1 - \frac{1}{2} \end{matrix} \right] (\kappa w, \tau + 1) &= e^{-\pi i (\alpha_1^2 + \alpha_1)} \vartheta \left[\begin{matrix} \alpha_1 \\ \beta_1 \end{matrix} \right] (\kappa w, \tau). \end{aligned}$$

Employing (84) it follows that

$$F \left[\begin{matrix} f \\ g \end{matrix} \right] \Big|_S (\widehat{x}) = \chi \left[\begin{matrix} f \\ g \end{matrix} \right] (S) F \left[\begin{matrix} f \\ g \end{matrix} \right] (\widehat{x}), \quad F \left[\begin{matrix} f \\ g \end{matrix} \right] \Big|_T (\widehat{x}) = \chi \left[\begin{matrix} f \\ g \end{matrix} \right] (T) F \left[\begin{matrix} f \\ g \end{matrix} \right] (\widehat{x}),$$

for S, T multipliers

$$\chi \left[\begin{matrix} f \\ g \end{matrix} \right] (S) = e^{-2\pi i \alpha_1 \beta_1}, \quad \chi \left[\begin{matrix} f \\ g \end{matrix} \right] (T) = e^{-\pi i (\alpha_1^2 + \alpha_1 + \frac{1}{12})}.$$

Thus it follows that $\widehat{Z}_{V_{\mathbb{Z}}}^{(2)} \left[\begin{matrix} f \\ g \end{matrix} \right] (\widehat{x})$ is invariant under all $\gamma_1 \in \Gamma_1$ up to a multiplier $\chi \left[\begin{matrix} f \\ g \end{matrix} \right] (\gamma_1)$ generated by S, T .

We next consider the action of the Heisenberg group \widehat{H} . From the definition of the theta series and using (83) we obtain

$$\begin{aligned} \vartheta \left[\begin{matrix} \alpha_1 - a\kappa \\ \beta_1 - b\kappa \end{matrix} \right] (\kappa(w + 2\pi i a\tau + 2\pi i b), \tau) &= e^{-(\pi i \tau a^2 + a w) \kappa^2} e^{-2\pi i a \kappa \beta_1} \vartheta \left[\begin{matrix} \alpha_1 \\ \beta_1 \end{matrix} \right] (\kappa w, \tau), \\ \exp \left(\frac{1}{2} \kappa^2 \widehat{l}(\mu(a, b, c) \cdot \widehat{x}) \right) &= e^{(\pi i \tau a^2 + a w + \pi i a b + \pi i c) \kappa^2} \exp \left(\frac{1}{2} \kappa^2 \widehat{l}(\widehat{x}) \right). \end{aligned}$$

³We note that the action quoted for the C generator in eqn (112) of [TZ1] contains an erroneous minus sign

Furthermore, from (85) we note that

$$\mu(a, b, c) \cdot \beta_2 = \beta_2 + a\beta_1 - b\alpha_1 - c\kappa + \frac{1}{2}(ab - c).$$

These results imply that

$$F \begin{bmatrix} f \\ g \end{bmatrix} \Big|_{\mu(a,b,c)} (\hat{x}) = \chi \begin{bmatrix} f \\ g \end{bmatrix} (\mu(a, b, c)) F \begin{bmatrix} f \\ g \end{bmatrix} (\hat{x}),$$

for multiplier

$$\chi \begin{bmatrix} f \\ g \end{bmatrix} (\mu(a, b, c)) = e^{-2\pi i b \alpha_1 \kappa} e^{\pi i (ab - c) \kappa (\kappa + 1)}.$$

Thus, altogether we have shown that

Theorem 5.10 $\widehat{Z}_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\hat{x})$ is holomorphic on $\widehat{\mathcal{D}}^\rho$ and is modular invariant with respect to $\gamma \in L$ with a multiplier system:

$$\widehat{Z}_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} \Big|_{\gamma} (\hat{x}) = \chi \begin{bmatrix} f \\ g \end{bmatrix} (\gamma) \widehat{Z}_{V_{\mathbb{Z}}}^{(2)} \begin{bmatrix} f \\ g \end{bmatrix} (\hat{x}).$$

These modular properties and multiplier system can also be confirmed by comparison to the modular properties of the genus two Riemann theta function expression of Theorem 5.8 using results of [MT2]. In particular, the multiplier $\chi \begin{bmatrix} f \\ g \end{bmatrix} (\mu(a, b, c))$ arises from the genus two theta function and can be easily independently verified.

Finally, in [TZ2] we showed that the Szegő kernel is modular invariant under L so that we conclude that the genus two generating function $\mathcal{G}_n^{(2)} \begin{bmatrix} f \\ g \end{bmatrix}$ lifts to a modular invariant form on $\widehat{\mathcal{D}}^\rho$ with the same multiplier.

6 Appendix

6.1 Proof of Proposition 3.3

Consider the coefficient of $z_0^{-N-1+\alpha\beta}$ of the Jacobi identity (37)

$$z_0^{-1} \left(\frac{z_1 - z_2}{z_0} \right)^{-\alpha\beta} \delta \left(\frac{z_1 - z_2}{z_0} \right) = \sum_{r \in \mathbb{Z}} (z_1 - z_2)^{r-\alpha\beta} z_0^{-r-1+\alpha\beta}.$$

Thus the coefficient of $z_0^{-N-1+\alpha\beta}$ arising from the first term on the left hand side of (37) is

$$(z_1 - z_2)^{N-\alpha\beta} \mathcal{Y}(u \otimes e^\alpha, z_1) \mathcal{Y}(v \otimes e^\beta, z_2).$$

A corresponding term arises from the second term on the left hand side of (37). On the other hand, by lower truncation, only finitely many negative integer powers of z_0 occur in

$$z_0^{-\alpha\beta} \mathcal{Y}(u \otimes e^\alpha, z_0) (v \otimes e^\beta),$$

so that (38) holds for $N \gg 0$.

Apply the operators of (38) to the vacuum vector together with translation and creativity to find

$$\begin{aligned} & (z_1 - z_2)^N (z_1 - z_2)^{-\alpha\beta} e^{z_2 L(-1)} \mathcal{Y}(u \otimes e^\alpha, z_1 - z_2) v \otimes e^\beta \\ = & (z_1 - z_2)^N (z_2 - z_1)^{-\alpha\beta} e^{z_1 L(-1)} \mathcal{Y}(v \otimes e^\beta, z_2 - z_1) u \otimes e^\alpha. \end{aligned}$$

Hence (39) follows for $z = z_1 - z_2$.

To prove (40) we consider the action on $w \otimes e^\gamma$ of the vertex operators appearing in the generalized Jacobi identity (37). The first term on the left hand side of (37) gives

$$\mathcal{Y}(u \otimes e^\alpha, z_1) \mathcal{Y}(v \otimes e^\beta, z_2) w \otimes e^\gamma = z_1^{\alpha(\beta+\gamma)} z_2^{\beta\gamma} A(z_1) B(z_2) w \otimes e^{\alpha+\beta+\gamma}, \quad (87)$$

for $A(z_1) = Y_-(\alpha, z_1) Y(u, z_1) Y_+(\alpha, z_1)$ and $B(z_2) = Y_-(\beta, z_2) Y(v, z_2) Y_+(\beta, z_2)$. We note that $A(z_1) B(z_2) w \otimes e^{\alpha+\beta+\gamma} = \sum_{k \in \mathbb{Z}} A(k) z_1^k B(z_2) w \otimes e^{\alpha+\beta+\gamma}$ for some operators $A(k)$. Furthermore

$$\begin{aligned} z_0^{-1} \left(\frac{z_1 - z_2}{z_0} \right)^{-\alpha\beta} \delta \left(\frac{z_1 - z_2}{z_0} \right) z_1^{k+\alpha\beta} &= \sum_{r \in \mathbb{Z}} \sum_{s \geq 0} \binom{r - \alpha\beta}{s} (-z_2)^s z_0^{-r+\alpha\beta-1} z_1^{r+k-s} \\ &= \sum_{N \in \mathbb{Z}} z_1^{-N-1} (z_0 + z_2)^{N+k+\alpha\beta}. \end{aligned}$$

Thus the coefficient of $z_1^{-N-1+\alpha\gamma}$ of (87) is

$$\begin{aligned} & (z_0 + z_2)^{N+\alpha\beta} z_2^{\beta\gamma} A(z_0 + z_2) B(z_2) w \otimes e^{\alpha+\beta+\gamma} \\ = & (z_0 + z_2)^{N-\alpha\gamma} \mathcal{Y}(u \otimes e^\alpha, z_0 + z_2) \mathcal{Y}(v \otimes e^\beta, z_2) w \otimes e^\gamma. \end{aligned}$$

On the other hand, the second term on left hand side of the Jacobi identity gives

$$\begin{aligned} & -z_0^{-1} \left(\frac{z_2 - z_1}{z_0} \right)^{-\alpha\beta} \delta \left(\frac{z_2 - z_1}{-z_0} \right) \mathcal{Y}(v \otimes e^\beta, z_2) \mathcal{Y}(u \otimes e^\alpha, z_1) w \otimes e^\gamma \\ & = -z_0^{-1} \left(\frac{z_2 - z_1}{z_0} \right)^{-\alpha\beta} \delta \left(\frac{z_2 - z_1}{-z_0} \right) z_1^{\alpha\gamma} z_2^{\beta(\alpha+\gamma)} B(z_2)A(z_1) w \otimes e^{\alpha+\beta+\gamma}. \end{aligned}$$

The coefficient of $z_1^{-N-1+\alpha\gamma}$ of this expression vanishes for $N \gg 0$ by lower truncation.

The right hand side of the Jacobi relation gives

$$\begin{aligned} & z_2^{-1} \delta \left(\frac{z_1 - z_0}{z_2} \right) \mathcal{Y}(\mathcal{Y}(u \otimes e^\alpha, z_0)(v \otimes e^\beta), z_2) \left(\frac{z_1 - z_0}{z_2} \right)^{\alpha\alpha(0)} w \otimes e^\gamma \\ & = z_2^{-1} \left(\frac{z_1 - z_0}{z_2} \right)^{\alpha\gamma} \delta \left(\frac{z_1 - z_0}{z_2} \right) \mathcal{Y}(\mathcal{Y}(u \otimes e^\alpha, z_0)(v \otimes e^\beta), z_2) w \otimes e^\gamma. \end{aligned}$$

But

$$\begin{aligned} z_2^{-1} \left(\frac{z_1 - z_0}{z_2} \right)^{\alpha\gamma} \delta \left(\frac{z_1 - z_0}{z_2} \right) & = \sum_{r \in \mathbb{Z}} \sum_{s \geq 0} \binom{r + \alpha\gamma}{s} (-z_0)^s z_1^{r-s+\alpha\gamma} z_2^{-r-\alpha\gamma-1} \\ & = \sum_{N \in \mathbb{Z}} z_1^{-N-1+\alpha\gamma} (z_2 + z_0)^{N-\alpha\gamma}. \end{aligned}$$

Taking the coefficient of $z_1^{-N-1+\alpha\gamma}$ of this expression leads to the result. Finally, (41) follows in the usual way on substituting $\alpha = 0$ into (37). \square

6.2 Proof of Proposition 4.1

We first show that the general intertwining n -point function can be written in terms of a 1-point function:

$$Z_\alpha^{(1)}(e^{\beta_1}, z_1; \dots; e^{\beta_n}, z_n; q) = Z_\alpha^{(1)}(\mathcal{Y}[e^{\beta_1}, z_{1n}] \dots \mathcal{Y}[e^{\beta_{n-1}}, z_{n-1n}] e^{\beta_n}; q) \quad (88)$$

$$= Z_\alpha^{(1)}(\mathcal{Y}[e^{\beta_1}, z_1] \dots \mathcal{Y}[e^{\beta_n}, z_n] \mathbf{1}; q), \quad (89)$$

where $\mathcal{Y}[e^{\beta_1}, z_1] \dots \mathcal{Y}[e^{\beta_n}, z_n] \mathbf{1} \in M[[z_i, z_i^{-1}]]$ since $\sum_{i=1}^n \beta_i = 0$. Using (32), (33) and the identity [TZ3]

$$e^{-\alpha\hat{q}} \mathcal{Y}(u \otimes e^\beta, z) e^{\alpha\hat{q}} = \mathcal{Y}(\Delta(\alpha, z) u \otimes e^\beta, z), \quad (90)$$

together with $\Delta(\alpha, q_i)e^{\beta_i} = q_i^{\alpha a(0)}e^{\beta_i}$ we find

$$\begin{aligned} & Z_\alpha^{(1)}(e^{\beta_1}, z_1; \dots; e^{\beta_n}, z_n; q) \\ &= \text{Tr}_M \left(e^{-\alpha \widehat{q}} \mathcal{Y} \left(q_1^{L(0)} e^{\beta_1}, q_1 \right) \dots \mathcal{Y} \left(q_n^{L(0)} e^{\beta_n}, q_n \right) q^{L(0)-1/24} e^{\alpha \widehat{q}} \right) \\ &= \text{Tr}_M \left(\mathcal{Y} \left(q_1^{L(0)+\alpha a(0)} e^{\beta_1}, q_1 \right) \dots \mathcal{Y} \left(q_n^{L(0)+\alpha a(0)} e^{\beta_n}, q_n \right) q^{L_g(0)-1/24} \right). \end{aligned}$$

Define the ‘‘shifted’’ conformal vector $\omega_\alpha = \omega - \alpha a(-2)\mathbf{1}$ with modes [DM]

$$L_\alpha(n) = L(n) + \alpha(n+1)a(n),$$

which satisfy the Virasoro algebra with central charge $c_\alpha = 1 - 12\alpha^2$ so that $(\mathcal{M}, \mathcal{Y}, \mathbf{1}, \omega_\alpha)$ is a Generalized VOA of central charge c_α . Noting that

$$L_g(0) - \frac{1}{24} = L_\alpha(0) - \frac{c_\alpha}{24},$$

we thus obtain (cf. Proposition 9 of [MTZ])

$$Z_\alpha^{(1)}(e^{\beta_1}, z_1; \dots; e^{\beta_n}, z_n; q) = \text{Tr}_M \left(\mathcal{Y} \left(q_1^{L_\alpha(0)} e^{\beta_1}, q_1 \right) \dots \mathcal{Y} \left(q_n^{L_\alpha(0)} e^{\beta_n}, q_n \right) q^{L_\alpha(0)-c_\alpha/24} \right).$$

Since we are tracing over M , we may consecutively apply the standard form of associativity to obtain:

$$\begin{aligned} & \text{Tr}_M \left(Y \left(\mathcal{Y} \left(q_1^{L_\alpha(0)} e^{\beta_1}, q_1 - q_n \right) \mathcal{Y} \left(q_2^{L_\alpha(0)} e^{\beta_2}, q_2 - q_n \right) \dots \right. \right. \\ & \quad \left. \left. \dots \mathcal{Y} \left(q_{n-1}^{L_\alpha(0)} e^{\beta_{n-1}}, q_n - q_n \right) q_n^{L_\alpha(0)} e^{\beta_n}, q_n \right) q^{L_\alpha(0)-c_\alpha/24} \right) \\ &= \text{Tr}_M \left(Y \left(q_n^{L_\alpha(0)} \mathcal{Y} [e^{\beta_1}, z_{1n}]_\alpha \dots \mathcal{Y} [e^{\beta_{n-1}}, z_{n-1n}]_\alpha e^{\beta_n}, q_n \right) q^{L_\alpha(0)-c_\alpha/24} \right), \quad (91) \end{aligned}$$

on using

$$q_n^{-L_\alpha(0)} \mathcal{Y}(e^\beta, q_i) q_n^{L_\alpha(0)} = \mathcal{Y}(q_n^{-L_\alpha(0)} e^\beta, q_n^{-1} q_i),$$

and where

$$\mathcal{Y} [e^\beta, z]_\alpha = \mathcal{Y} (q_z^{L_\alpha(0)} e^\beta, q_z - 1),$$

denotes the square bracket vertex operator for the shifted Virasoro system.

But the identities

$$\begin{aligned} \Delta(-\alpha, q_n) q_n^{L_\alpha(0)} &= q_n^{L(0)} \Delta(-\alpha, 1), \\ \Delta(-\alpha, 1) \mathcal{Y} [e^\beta, z]_\alpha \Delta(\alpha, 1) &= \mathcal{Y} [e^\beta, z], \end{aligned}$$

imply that (91) becomes the 1-point function (88). (89) follows since

$$\begin{aligned} o(\mathcal{Y}[e^{\beta_1}, z_1] \dots \mathcal{Y}[e^{\beta_n}, z_n] \mathbf{1}) &= o(\mathcal{Y}[e^{\beta_1}, z_1] \dots \mathcal{Y}[e^{\beta_{n-1}}, z_{n-1}] e^{z_n L[-1]} e^{\beta_n}) \\ &= o(e^{z_n L[-1]} \mathcal{Y}[e^{\beta_1}, z_{1n}] \dots \mathcal{Y}[e^{\beta_{n-1}}, z_{n-1n}] e^{\beta_n}) \\ &= o(\mathcal{Y}[e^{\beta_1}, z_{1n}] \dots \mathcal{Y}[e^{\beta_{n-1}}, z_{n-1n}] e^{\beta_n}), \end{aligned}$$

and since $o(L[-1]u) = 0$ for all $u \in M$ [Z1].

As in the proof of Proposition 5 of [MT1] using the identity

$$Y_+(\alpha, x)Y_-(\beta, y) = \left(1 - \frac{y}{x}\right)^{\alpha\beta} Y_-(\beta, y)Y_+(\alpha, x),$$

we find that (89) is given by

$$\prod_{1 \leq r < s \leq n} z_{rs}^{\beta_r \beta_s} Z_\alpha^{(1)} \left(\exp \left(\sum_{m>0} \frac{a[-m]}{m} \sum_{i=1}^n \beta_i z_i^m \right) \mathbf{1}; q \right).$$

Proposition 3 of [MT1] gives

$$\begin{aligned} &Z_\alpha^{(1)} \left(\exp \left(\sum_{m>0} \frac{a[-m]}{m} \sum_{i=1}^n \beta_i z_i^m \right) \mathbf{1}; q \right) \\ &= q^{\frac{1}{2}\alpha^2} \exp \left(\alpha \sum_{i=1}^n \beta_i z_i \right) Z_0 \left(\exp \left(\sum_{m>0} \frac{a[-m]}{m} \sum_{i=1}^n \beta_i z_i^m \right) \mathbf{1}; q \right). \end{aligned}$$

Finally, Proposition 4 of [MT1] gives

$$Z_0 \left(\exp \left(\sum_{m>0} \frac{a[-m]}{m} \sum_{i=1}^n \beta_i z_i^m \right) \mathbf{1}; q \right) = \frac{1}{\eta(q)} \prod_{1 \leq r < s \leq n} \left(\frac{K(z_{rs}, \tau)}{z_{rs}} \right)^{\beta_r \beta_s}.$$

The proposition follows. \square

References

- [BK] Bakalov, B. and Kac, V., *Generalized Vertex Algebras*, math.QA/0602072 .
- [DL] Dong, C. and Lepowsky, J.: *Generalized Vertex Algebras and Relative Vertex Operators*, Progress in Mathematics **112** Birkhäuser (Boston, MA, 1993).

- [DLinM] Dong, C., Lin, Z., Mason, G.: On vertex operator algebras as sl_2 -modules. Arasu, K.T. (ed.) et al., *Groups, difference sets, and the Monster*, Proceedings of a special research quarter, Columbus, OH, USA, Spring 1993, Walter de Gruyter, (Berlin 1996), Ohio State Univ. Math. Res. Inst. Publ. **4** (1996) 349–362.
- [DM] Dong, C. and Mason, G.: Shifted vertex operator algebras, *Math. Proc. Camb. Philos. Soc.* **141** (2006) 67–80.
- [DP] D’Hoker, E. and Phong, D.H., The geometry of string perturbation theory, *Rev. Mod. Phys.* **60** (1988) 917–1065.
- [DVPFHLS] Di Vecchia, P., Pezzella, F., Frau, M., Hornfeck, K., Lerda, A. and Sciuto, S.: N -point g -loop vertex for a free fermionic theory with arbitrary spin. *Nucl. Phys.* **B333** (1990) 635–700.
- [EO] Eguchi, T. and Ooguri, H.: Conformal and current algebras on a general Riemann surface, *Nucl. Phys.* **B282** (1987) 308–328.
- [EZ] Eichler, M. and Zagier, D.: *The Theory of Jacobi Forms*, Birkhäuser (Boston, 1985).
- [F] Fay, J.D.: *Theta Functions on Riemann Surfaces*, Lecture Notes in Mathematics, Springer-Verlag, (Berlin-New York, 1973).
- [FHL] Frenkel, I., Huang, Y. and Lepowsky, J.: On axiomatic approaches to vertex operator algebras and modules, *Mem. Amer. Math. Soc.* **104** (1993).
- [FK] Farkas, H.M. and Kra, I., *Theta Constants, Riemann surfaces and the Modular Group*, Graduate Studies in Mathematics, **37**, AMS (Providence, 2001).
- [FLM] Frenkel, I., Lepowsky, J. and Meurman, A., *Vertex Operator Algebras and the Monster*, Academic Press, (New York, 1988).
- [HS] Hawley, N.S. and Schiffer, M.: Half-order differentials on Riemann surfaces, *Acta Math.* **115** (1966) 199–236.

- [K] Kac, V.: *Vertex Operator Algebras for Beginners*, University Lecture Series **10**, AMS (Providence 1998).
- [Li] Li, H.: Symmetric invariant bilinear forms on vertex operator algebras, *J. Pure. Appl. Alg.* **96** (1994) 2790–297.
- [M] Mumford, D.: *Tata lectures on Theta I and II*, Birkhäuser, (Boston, 1983).
- [MN] Matsuo, A. and Nagatomo, K.: Axioms for a vertex algebra and the locality of quantum fields, *Math. Soc. Jap. Mem.* **4** (1999).
- [MT1] Mason, G. and Tuite, M.P.: Free bosonic vertex operator algebras on genus two Riemann surfaces I, *Commun. Math. Phys.* **300** (2010) 673–713.
- [MT2] Mason, G. and Tuite, M.P.: On genus two Riemann surfaces formed from sewn tori, *Commun. Math. Phys.* **270** (2007) 587–634.
- [MT3] Mason, G. and Tuite, M.P.: Free bosonic vertex operator algebras on genus two Riemann surfaces II, arXiv:1111.2264, to appear in *Proceedings of Conformal Field Theory, Automorphic Forms and Related Topics*, Heidelberg, 2011.
- [MT4] Mason, G. and Tuite, M.P.: Chiral n -point functions for free boson and lattice vertex operator algebras, *Commun. Math. Phys.* **235** (2003) 47–68.
- [MT5] Mason, G. and Tuite, M.P.: Vertex operators and modular forms, *A Window into Zeta and Modular Physics* eds. K. Kirsten and F. Williams, MSRI Publications **57** 183–278, Cambridge University Press, (Cambridge, 2010).
- [MTZ] Mason, G., Tuite, M.P. and Zuevsky, A.: Torus n -point functions for \mathbb{R} -graded vertex operator superalgebras and continuous fermion orbifolds, *Commun. Math. Phys.* **283** (2008) 305–342.
- [R] Raina, A.K.: Fay’s trisecant identity and conformal field theory, *Comm. Math. Phys.* **122**, (1989) 625–641.

- [S] Scheithauer, N.: Vertex algebras, Lie algebras and superstrings, *J. Alg.* **200** (1998) 363–403.
- [T] Tuite, M.P: Genus two meromorphic conformal field theory, *CRM Proceedings and Lecture Notes* **30** (2001) 231–251.
- [TUY] Tsuchiya, A., Ueno, K. and Yamada, Y.: Conformal field theory on universal family of stable curves with gauge symmetries, *Adv. Stud. Pure. Math.* **19** (1989) 459–566.
- [TZ1] Tuite, M.P. and Zuevsky, A.: The Szegő kernel on a sewn Riemann surface, *Commun. Math. Phys.*, **306**, (2011) 617–645.
- [TZ2] Tuite, M.P. and Zuevsky, A.: Genus two partition and correlation functions for fermionic vertex operator superalgebras I, *Comm. Math. Phys.*, **306**, (2011) 419–447.
- [TZ3] Tuite, M.P. and Zuevsky, A.: A generalized vertex operator algebra for Heisenberg intertwiners, *J. Pure Appl. Alg.* **216** (2012) 1442–1453.
- [Z1] Zhu, Y.: Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996) 237–302.
- [Z2] Zhu, Y.: Global vertex operators on Riemann surfaces, *Commun. Math. Phys.* **165** (1994) 485–531.