




## Impaired endocannabinoid signalling in the rostral ventromedial medulla underpins genotype-dependent hyper-responsivity to noxious stimuli

Title	Impaired endocannabinoid signalling in the rostral ventromedial medulla underpins genotype-dependent hyper-responsivity to noxious stimuli
Author(s)	Rea, Kieran;Olango, Weredeselam M.;Okine, Bright N.;Madasu, Manish K.;McGuire, Iseult C.;Coyle, Kathleen;Harhen, Brendan;Roche, Michelle;Finn, David P.
Publication Date	2014-01
Publisher	International Association for the Study of Pain
Repository DOI	<a href="https://doi.org/10.1016/j.pain.2013.09.012">10.1016/j.pain.2013.09.012</a>

**AUTHOR QUERY FORM**

	<b>Journal: PAIN</b>  <b>Article Number: 8989</b>	<b>Please e-mail or fax your responses and any corrections to:</b>  <b>E-mail: <a href="mailto:proofcorrections.eseo@elsevier.sps.co.in">proofcorrections.eseo@elsevier.sps.co.in</a></b>  <b>Fax: +353-61-709240</b>
---	---	---

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

<b>Location in article</b>	<b>Query / Remark: <a href="#">click on the Q link to go</a> Please insert your reply or correction at the corresponding line in the proof</b>
<a href="#">Q1</a>	Please confirm that given name(s) and surname(s) have been identified correctly.
<a href="#">Q2</a>	Please advise whether "Department of" may be inserted at the beginning of each affiliation.
<a href="#">Q3</a>	Please approve Summary as edited.
<a href="#">Q4</a>	Please expand "CB <sub>1</sub> " (cannabinoid-1?) at first mention in abstract and text.
<a href="#">Q5</a>	Please provide city for Harlan and any other manufacturers for which this information has not been provided.
<a href="#">Q6</a>	Please verify that "25% v/v 1 mol/L Tris HCl, pH 6.8," is correct ("1M" changed to "1 mol/L").
<a href="#">Q7</a>	Please correct fifth author surname in Ref. [13] (please note that it has become corrupted during file conversion).
<a href="#">Q8</a>	Please expand "AAPSJ" to the standard abbreviation in Refs. [39] and [75].
<a href="#">Q9</a>	Please provide city and state for publisher in Ref. [66].
<a href="#">Q10</a>	Please provide journal volume and verify article page range in Ref. 88.
	<div style="border: 1px solid black; padding: 5px; margin-top: 20px;"> <p style="color: red;">Please check this box if you have no corrections to make to the PDF file</p> <input style="width: 40px; height: 20px; vertical-align: middle;" type="checkbox"/> </div>

Thank you for your assistance.

### Summary

---

#### **Impaired endocannabinoid signalling in the rostroventromedial medulla underpins genotype-dependent hyper-responsivity to noxious stimuli**

- Q3 Impaired endocannabinoid signalling in the [rostroventromedial](#) medulla underpins hyper-responsivity to a noxious inflammatory stimulus in the Wistar-Kyoto rat, a genetic background prone to heightened stress/affect.





PAIN® xxx (2013) xxx–xxx

PAIN®

www.elsevier.com/locate/pain

# Impaired endocannabinoid signalling in the rostromedial medulla underpins genotype-dependent hyper-responsivity to noxious stimuli

Kieran Rea<sup>a,c,1</sup>, Weredeslam M. Olango<sup>a,c,1</sup>, Bright N. Okine<sup>a,c</sup>, Manish K. Madasu<sup>a,c</sup>, Iseult C. McGuire<sup>a,c</sup>,  
Cathleen C. Brennan<sup>c</sup>, Brendan Harhen<sup>c</sup>, Michelle Roche<sup>b,c</sup>, David Finn<sup>a,b,\*</sup>

<sup>a</sup>Pharmacology and Therapeutics, National University of Ireland, Galway, Ireland  
<sup>b</sup>Physiology, School of Medicine, National University of Ireland, Galway, Ireland  
<sup>c</sup>NCBES Neuroscience Cluster and Centre for Pain Research, National University of Ireland, Galway, Ireland

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

## ARTICLE INFO

**Article history:**  
Received 28 April 2013  
Received in revised form 18 August 2013  
Accepted 9 September 2013  
Available online xxxxx

**Keywords:**  
Pain  
Affect  
Wistar-Kyoto rat  
Cannabinoid<sub>1</sub> (CB<sub>1</sub>) receptor  
Anandamide  
Fatty acid amide hydrolase (FAAH)  
Formalin  
Rostromedial medulla (RVM)

## ABSTRACT

Pain is both a sensory and an emotional experience, and is subject to modulation by a number of factors including genetic background modulating stress/affect. The Wistar-Kyoto (WKY) rat exhibits a stress-hyperresponsive and depressive-like phenotype and increased sensitivity to noxious stimuli, compared with other rat strains. Here, we show that this genotype-dependent hyperalgesia is associated with impaired pain-related mobilisation of endocannabinoids and transcription of their synthesising enzymes in the rostromedial medulla (RVM). Pharmacological blockade of the CB<sub>1</sub> receptor potentiates the hyperalgesia in WKY rats, whereas inhibition of the endocannabinoid catabolising enzyme, fatty acid amide hydrolase, attenuates the hyperalgesia. The latter effect is mediated by CB<sub>1</sub> receptors in the RVM. Together, these behavioural, neurochemical, and molecular data indicate that impaired endocannabinoid signalling in the RVM underpins hyper-responsivity to noxious stimuli in a genetic background prone to heightened stress/affect.

© 2013 International Association for the Study of Pain. Published by Elsevier B.V. All rights reserved.

## 1. Introduction

The ability to perceive pain and respond appropriately is essential for survival. However, excessive or persistent pain constitutes a major healthcare problem for those who experience it, and for society and economies. Pain is both a sensory and an emotional experience and is subject to modulation by a number of factors. A key factor is the contribution of genetic background and its influence on stress responding and affective processing. An increased understanding of how such factors can influence pain is important from a fundamental physiological perspective, and may also aid the identification of new therapeutic targets for the treatment of persistent pain and its exacerbation by, and/or co-morbidity with, stress-related affective disorders.

The influence of genetic background and stress/affect on pain can be examined by comparing behavioural, neurochemical, and molecular responses to noxious stimuli across different rodent

strains. The Wistar-Kyoto (WKY) inbred rat strain exhibits a stress-hyperresponsive and depressive-like phenotype [5,29,63,64] and displays increased sensitivity to visceral and somatic noxious stimuli, compared with other rat strains [5,31,33,71,85,90]. As such, the WKY rat represents a useful model with which to study the impact of genetic background and negative affect on pain processing.

The endogenous cannabinoid (endocannabinoid) system plays a key role in the modulation of both pain processing and emotionality [23,27,38,39,48,87,88]. This system comprises at least 2 receptors, the CB<sub>1</sub> [14,52] and CB<sub>2</sub> [58] receptors, of which the CB<sub>1</sub> receptor is most abundant in the brain. *N*-arachidonylethanolamide (anandamide: AEA [15]) and 2-arachidonoylglycerol (2-AG [53,83]) are the 2 most extensively studied endogenous ligands for the cannabinoid receptors. AEA and 2-AG are synthesized from phospholipid precursors by *N*-acyl phosphatidylethanolamine phospholipase D (NAPE-PLD) [17,45] and diacylglycerol lipase (DAGL) [75], respectively, and are catabolised primarily by fatty acid amide hydrolase (FAAH) [10,62,76] and monoacylglycerol lipase [MAGL] [30], respectively. Both endocannabinoids have similar affinity for both CB<sub>1</sub> [47] and CB<sub>2</sub> [32] receptors; however, 2-AG usually has higher efficacy than AEA at both receptors [32,47]. Work from our laboratory and others has demonstrated

\* Corresponding author. Address: Pharmacology and Therapeutics, School of Medicine, University Road, National University of Ireland, Galway, Ireland. Tel.: +353 91 495280; fax: +353 91 495586.

E-mail address: david.finn@nuigalway.ie (D.P. Finn).

<sup>1</sup> K.R. and W.M.O. contributed equally to this work.

an important role for these endocannabinoids in stress–pain interactions, regulating both stress-induced analgesia [6–8,28,60,67,69] and stress-induced hyperalgesia [40,77]. To date, however, the contribution of the endocannabinoid system to altered nociceptive responding in genetic backgrounds predisposed to negative affect has not been investigated. Research has shown that the endocannabinoid system of WKY rats is dysfunctional, with altered expression of endocannabinoid catabolising enzymes likely contributing to their depressive phenotype [86]. However, the extent to which alterations in the endocannabinoid system may explain altered nociceptive responding in WKY rats is unknown, and was the focus of the current studies.

We have used behavioural, neurochemical, and molecular approaches to test the hypothesis that enhanced pain-related behavioural responding to the noxious inflammatory stimulus of intra-plantar formalin injection in WKY rats is mediated by impaired mobilisation of endocannabinoid-CB<sub>1</sub> receptor signalling. Our studies focused on the role of the endocannabinoid system in the rostromedial medulla (RVM), given its key role in top–down descending modulation of pain [21,22,36,91], and evidence that CB<sub>1</sub> receptors in the RVM [37,51] regulate nociceptive processing [50,55,84].

## 2. Methods

### 2.1. Animals

For all experiments, male Sprague–Dawley (SD) and/or Wistar–Kyoto (WKY) rats (Harlan, UK) were used. Animals were singly housed, and holding rooms were maintained at a constant temperature (21 ± 2°C) under standard lighting conditions (12:12-hour light/dark, lights on from 0800 to 2000 h). Experiments were carried out during the light phase between 0800 and 1700 h. Food and water were available ad libitum. The experiments adhered to the guidelines of the Committee for Research and Ethical Issues of IASP [[www.iasp-pain.org/AM/Template.cfm?Section=Animal\\_Research](http://www.iasp-pain.org/AM/Template.cfm?Section=Animal_Research)]. The experimental protocol was carried out after approval by the Animal Care and Research Ethics Committee, National University of Ireland, Galway, under license from the Irish Department of Health and Children and in compliance with the European Communities Council directive 86/609.

### 2.2. Experimental design

Three separate experiments were performed. In all experiments, animals were randomly assigned to treatment groups, and the sequence of treatments and testing was also randomised to control for the order of testing. Experiment 1 investigated whether enhanced formalin-evoked nociceptive behaviour in WKY rats vs their SD counterparts was associated with alterations in endocannabinoid levels in the RVM or genes coding for the enzymes and receptors of the endocannabinoid system. A total of 24 male Sprague–Dawley rats and 24 male WKY rats (285–320 g) received an intra-plantar injection of 50 µL formalin (2.5% in 0.9% saline, s.c.) or 0.9% saline (control group) into the right hindpaw immediately after a 10-minute habituation exposure to the formalin test arena. This design resulted in 4 experimental groups, as follows: SD-Saline (SD-Sal); SD-Formalin (SD-Form); WKY-Saline (WKY-Sal); and WKY-Formalin (WKY-Form) (n = 10–12 per group). At the peak of the second phase of the formalin test (30 minutes after formalin injection), rats were killed by decapitation. Brains were removed rapidly and were snap-frozen on dry ice and stored at –80°C before microdissection of the RVM and subsequent analysis of AEA and 2-AG levels using liquid chromatography coupled to tandem mass spectrometry

(LC-MS/MS). Microdissected RVM tissue was also analysed by quantitative reverse transcription–polymerase chain reaction (RT-PCR) for expression of genes coding for the CB<sub>1</sub> receptor and for the endocannabinoid-related enzymes NAPE-PLD, DAGLα, FAAH, and MAGL. A separate cohort of rats (n = 6 per group) were treated exactly as described above to generate RVM tissue for western blot analysis of CB<sub>1</sub> receptor expression.

In experiment 2, we investigated the effects of pharmacological blockade of the CB<sub>1</sub> receptor or inhibition of FAAH on formalin-evoked nociceptive behaviour in WKY and SD rats. A total of 32 male Sprague–Dawley rats and 32 male Wistar–Kyoto rats (250–350 g) were assessed in the formalin test, with subjects receiving intraperitoneal (i.p.) injection of the CB<sub>1</sub> receptor antagonist/inverse agonist AM251 (3 mg/kg), the FAAH inhibitor URB597 (0.5 mg/kg) or vehicle (ethanol:cremaphor:saline vehicle in a ratio of 1:1:18; 3 mL/kg) before intra-plantar formalin injection. Rats were habituated to the formalin test arena for 10 minutes before formalin injection. URB597 and AM251 were administered 60 minutes and 30 minutes before formalin injection, respectively, based on previous studies in our laboratory and others demonstrating their *in vivo* efficacy at these doses and times of administration [1,7,8,34,35,41,44,49]. To control for the different times of injection of the 2 drugs, half of the vehicle-treated control rats received vehicle at 30 minutes and half at 60 minutes before intra-plantar formalin injection. These 2 vehicle-treated cohorts were subsequently combined as 1 group after statistical analysis confirmed that there were no differences between them for any of the experimental parameters examined. This design resulted in 6 experimental groups (n = 6–10 per group): SD-Vehicle [SD-Veh]; SD-AM251 (3 mg/kg) [SD-AM251]; SD-URB597 (0.5 mg/kg) [SD-URB]; WKY-Vehicle [WKY-Veh]; WKY-AM251 (3 mg/kg) [WKY-AM251]; and WKY-URB597 (0.5 mg/kg) [WKY-URB]. At the end of the formalin test (ie, 70 minutes after formalin injection), the rats were killed by decapitation.

Experiment 3 was conducted to investigate whether URB597-mediated suppression of formalin-induced hyperalgesia (result from experiment 2) in the WKY rats is mediated by AEA-induced activation of CB<sub>1</sub> receptors in the RVM. Male Wistar–Kyoto rats (280–350 g; Harlan, UK) were implanted with stainless steel guide cannulae targeting the RVM. On the test day, URB597 (0.5 mg/kg) or vehicle was administered by i.p. injection 60 minutes before formalin injection. Fifteen minutes before formalin injection, 0.3 µL of AM251 (1 µg/0.3 µL) or dimethylsulfoxide vehicle (DMSO, 100%) was microinjected over 1 minute through an injection needle that protruded 1 mm beyond the tip of the pre-implanted guide cannula, with the aid of a Hamilton microsyringe attached to polyethylene tubing and a Harvard PHD2000 infusion pump (Harvard Apparatus, Kent, UK) as described previously [28,60,67,69,72,73]. The injection needle was left in place for 1 minute more after infusion to allow for drug diffusion away from the injector tip. This dose of AM251 was chosen on the basis of previous studies demonstrating that microinjection of AM251 into different regions of the brain modulated behavioural responses to analgesic compounds in various animal models of pain [18,12,20]. After microinjection of AM251 or DMSO vehicle directly into the RVM, animals were immediately placed in a Perspex arena to habituate for 10 minutes. Animals were subsequently injected with formalin under brief anaesthesia and returned to the formalin test arena for behavioural analysis. They were killed by decapitation at 70 minutes after formalin administration. A 0.3-µL quantity of 1% fast green dye was microinjected via the guide cannula, and brains were rapidly removed, snap-frozen on dry ice, and stored at –80°C before injection site verification. Microdissection of the RVM was performed in conjunction with injection site verification, and the microdissected tissue was analysed for endocannabinoid levels using LC-MS/MS.

212 2.3. Drug preparation

213 The FAAH inhibitor URB597 [(3-(3-carbomoylphenyl)phenyl)N-  
214 cyclohexylcarbamate] and formalin were purchased from Sigma  
215 Aldrich (Dublin, Ireland). The CB<sub>1</sub> receptor antagonist/inverse ago-  
216 nist AM251 (N-(Piperidin-1-yl)-5-(4-iodophenyl)-1-(2,4-dichloro-  
217 phenyl)-4-methyl-1H-pyrazole-3-carboxamide was purchased  
218 from Abcam (Cambridge, UK). URB597 (0.5 mg/kg) or vehicle was  
219 administered by i.p. injection 60 minutes before formalin injection,  
220 whereas AM251 (3.0 mg/kg) was administered by i.p. injection  
221 30 minutes before formalin injection. Both drugs were reconsti-  
222 tuted as an emulsion in ethanol:cremaphor:saline vehicle in a ratio  
223 of 1:1:18 and administered at an injection volume of 3 mL/kg. For  
224 intra-RVM microinjections, AM251 was prepared to a concentra-  
225 tion of 1 µg per 0.3 µL of DMSO vehicle (dimethylsulfoxide,  
226 100%) and 0.3 µL was microinjected with the aid of a Hamilton  
227 microsyringe as described above.

228 2.4. Formalin test

229 Rats were placed in a Perspex observation chamber  
230 (30 × 30 × 40 cm; LxWxH) at 30 lux for a 10-minute habituation  
231 period, after which time they received an intra-plantar injection  
232 of 50 µL formalin (2.5% in 0.9% saline) or 0.9% saline into the right  
233 hindpaw under brief isoflurane anaesthesia as described previ-  
234 ously [7,8,24–26,28,67,69,72,73]. Rats were returned to their home  
235 cage for another 3 minutes, at which point they were returned to  
236 the same Perspex observation chamber to which they had previ-  
237 ously been exposed. A video camera located beneath the observa-  
238 tion chamber was used to record animal behaviour onto DVD for  
239 subsequent analysis. Behaviour was analysed with the aid of  
240 EthoVision XT7 software by a rater blinded to treatments. Formali-  
241 n-evoked nociceptive behaviour was categorized as time spent  
242 raising the formalin-injected paw above the floor without contact  
243 with any other surface (C1) and holding, licking, biting, shaking,  
244 or flinching the injected paw (C2) to obtain a composite pain score  
245 [CPS = (C1 + 2(C2))/(total duration of analysis period)] according to  
246 the method of Watson et al. [89].

247 2.5. Punch microdissection of RVM tissue

248 In experiments 1 and 3, frozen coronal brain sections (300-µm  
249 in thickness) containing the RVM were cut on a cryostat (MICROM,  
250 Germany). A series of 300-µm-thick sections (from AP -9.16 to  
251 -11.6 mm relative to bregma) were punched using cylindrical  
252 brain punchers (Harvard Apparatus; internal diameter 2 mm), with  
253 the aid of the rat brain atlas of Paxinos and Watson [66]. Punched  
254 RVM samples encompassed the gigantocellular reticularis nucleus,  
255 raphe magnus nucleus, medial lemniscus, raphe pallidus nucleus,  
256 pyramidal tracts, ventral aspect of the pontine reticular nucleus,  
257 and trigeminothalamic tract. These samples were weighed and  
258 stored at -80°C before extraction for determination of the concen-  
259 trations of the endocannabinoids by LC-MS/MS, gene expression  
260 analysis using quantitative RT-PCR or protein expression using  
261 western blotting.


262 2.6. Quantitative RT-PCR analysis of the expression of  
263 endocannabinoid-related genes

264 Total RNA was extracted from post-mortem tissue using a  
265 Machery–Nagel extraction kit (Nucleospin RNA II; Technopath, Ire-  
266 land) according to the manufacturer’s instructions and as previ-  
267 ously described [42,43]. RNA quality (1.8–2 260/280 ratio) and  
268 quantity was assessed using a Nanodrop spectrophotometer  
269 (ND-1000; Nanodrop, Labtech International, UK) and normalised  
270 to a concentration of 5 ng/µL. A 50-ng quantity of RNA from each

sample was reverse transcribed to cDNA using an Invitrogen  
Superscript III reverse transcriptase custom kit (Bio-Sciences, Dun  
Laoghaire, Ireland). Taqman gene expression assays (Applied  
Biosystems, UK) containing forward and reverse primers and a  
FAM-labelled MGB Taqman probe were used to quantify the gene  
of interest using an Applied Biosystems ‘stepOne plus’ instrument  
(Bio-Sciences, Dun Laoghaire, Ireland). Assay IDs for the genes  
examined were as follows for rat CB<sub>1</sub> (Rn00562880\_m1), FAAH  
(Rn00577086\_m1), MAGL (Rn00593297\_m1), NAPE-PLD  
(Rn01786262\_m1), and DAGL $\alpha$  (Rn01454304\_m1). VIC-labelled  
GAPDH (4308313) was used as the house-keeping gene and endog-  
enous control. A no-template control (NTC) reaction was included  
in all assays. The relative expression of target genes to endogenous  
control was calculated using the formula 2<sup>-ΔCt</sup>, where ΔCt repre-  
sents the magnitude of the difference between cycle threshold (Ct)  
values of the target and endogenous control, and the result was ex-  
pressed as a percentage of the mean value of the control group.

271 2.7. Western blot analysis of CB<sub>1</sub> receptor expression in the RVM 288

272 Western blotting was performed according to methods  
273 described previously [59], with minor modifications. Briefly, frozen  
274 punches of the RVM weighing approximately 10 mg were lysed by  
275 brief (3-second) sonication in radio-immunoprecipitation assay  
276 (RIPA) lysis buffer (150 mmol/L NaCl, 25 mmol/L Tris-HCl, pH 7.6,  
277 0.5% Triton X-100, 1% sodium deoxycholate, 0.1% sodium dodecyl  
278 sulphate, 1 mmol/L Na<sub>3</sub>VO<sub>4</sub>, 10 mmol/L NaF containing 1% protease  
279 inhibitor cocktail [Sigma-Aldrich, Dublin, Ireland]) at a ratio of 1:10  
280 (w/v) in a 1.5-mL microcentrifuge tube. Homogenate was placed  
281 on a shaker for 45 minutes at 4° with gentle agitation to allow  
282 for complete dissociation of nucleo-protein complexes and then  
283 centrifuged at 13,200 rpm (Eppendorf Centrifuge 5415R Stevenage,  
284 UK) for 20 minutes at 4°C. The supernatant was collected and  
285 protein content determined by Bradford assay [4]. A 36-µg quan-  
286 tity of protein sample in loading buffer (4X sample loading buffer:  
287 25% v/v 1 mol/L Tris HCl, pH 6.8, 5% w/v sodium dodecyl sulfate  
288 (SDS), 20% v/v glycerol, 2.5% Bromophenol blue (0.2% w/v in  
289 100% ethanol), and 20% v/v of 2-mercaptoethanol, made up to a  
290 total volume of 20 mL in distilled water), was boiled at 100°C for  
291 5 minutes, briefly centrifuged, and subjected to 9% SDS-polyacryl-  
292 amide gel electrophoresis (SDS-PAGE) at a constant voltage of  
293 120 mV for 2 hours. The separated protein samples were electro-  
294 blotted onto a nitrocellulose membrane (Nitrocellulose membrane,  
295 CAS# 9004-70-0; Bio-Rad, Dublin, Ireland) at 100 mV for 1 hour.  
296 Protein transfer efficiency was verified by ponceau (0.1% ponceau  
297 dye in 5% acetic acid) staining. Membranes were blocked in 5%  
298 non-fat dry milk in 0.05% Tris-buffered saline/Tween 20 (TBST)  
299 solution for 1 hour at room temperature and incubated with rabbit  
300 polyclonal antibody to the CB<sub>1</sub> receptor (C-term) (1:200, catalog  
301 no. 10006590; Cayman Chemical, MI) and mouse monoclonal anti-  
302 body to β-Actin (1:10,000, A5441; Sigma-Aldrich, Dublin, Ireland)  
303 diluted in 5% milk/0.05% TBST overnight at 4°C. Membranes were  
304 subjected to 3 10-minute washes in 0.05% TBST and incubated with  
305 secondary antibody solution containing IRDye conjugated goat  
306 anti-rabbit (λ800) and goat anti-mouse (λ700) (LI-COR Biosciences  
307 Abingdon Park, Oxford, UK) diluted 1:10,000 in 1% milk/0.05% TBST  
308 for 1 hour. Three 5-minute washing steps were then performed  
309 with washing solution and 1 final 5-minute wash in distilled  
310 H<sub>2</sub>O. Blots were scanned on a LI-COR Odyssey imager. IR band  
311 intensities for glycosylated (~62-kDa), and non-glycosylated  
312 (~53-kDa) CB<sub>1</sub> receptor protein expression [19] and β-actin  
313 (~42-kDa) for each sample were generated automatically using  
314 the background subtraction method of the LI-COR Image Studio  
315 Ver. 2.0 imaging software. The ratio of CB<sub>1</sub> receptor intensity to  
316 β-actin intensity was then calculated for each sample, and ex-  
317 pressed as a percentage of mean SD saline values. 334



**Fig. 1.** WKY rats display a greater nociceptive response to intra-plantar formalin administration compared with their SD counterparts over the total duration of the 30-minute trial. (a) Temporal profile of formalin-evoked nociceptive behaviour in SD and WKY rats. (b) Graphic representation of formalin-evoked nociceptive behaviour in SD and WKY rats over entire 30-minute trial. Two-way ANOVA (effects of strain:  $F_{1,45} = 10.436, P = .002$ ; formalin:  $F_{1,45} = 462.314, P < .001$  and strain  $\times$  formalin interaction:  $F_{1,45} = 10.347, P = .002$ ) followed by Fisher's LSD post-hoc test (### $P < .001$  vs SD-Sal, \*\*\* $P < .001$  vs WKY-Sal, \*\*\* $P < .001$  vs SD-Form). Data are expressed as mean  $\pm$  SEM ( $n = 10-12$  rats per group). Form, formalin; Sal, saline solution; SD, Sprague-Dawley; WKY, Wistar-Kyoto.

335 2.8. Quantitation of endocannabinoids in RVM tissue using LC-MS/MS


336 Quantitation of endocannabinoids was essentially as described  
 337 previously [7,28,42,43,60,68]. In brief, each sample was first  
 338 homogenized in 400  $\mu$ L 100% acetonitrile containing known fixed  
 339 amounts of deuterated internal standards (0.014 nmol AEA-d8  
 340 and 0.48 nmol 2-AG-d8). Homogenates were centrifuged at  
 341 14,000g for 15 minutes at 4°C, and the supernatant was collected  
 342 and evaporated to dryness. Lyophilized samples were resuspended  
 343 in 40  $\mu$ L 65% acetonitrile, and a 2- $\mu$ L quantity was injected onto a  
 344 Zorbax C18 column (150  $\times$  0.5 mm internal diameter) from a  
 345 cooled autosampler maintained at 4°C (Agilent Technologies, Cork,  
 346 Ireland). Mobile phases consisted of A (high-performance liquid  
 347 chromatography [HPLC]-grade water with 0.1% formic acid)  
 348 and B (acetonitrile with 0.1% formic acid), with a flow rate of  
 349 12  $\mu$ L/min. Reverse-phase gradient elution began initially at 65%  
 350 B and over 10 min was ramped linearly up to 100% B. At 10 min-  
 351 utes, the gradient was held at 100% B up to 20 minutes. At  
 352 20.1 minutes, the gradient returned to initial conditions for an-  
 353 other 10 minutes to re-equilibrate the column. Analyte detection  
 354 was carried out in electrospray-positive ionization mode on an Agi-  
 355 lent 1100 HPLC system coupled to a triple quadrupole 6460 mass  
 356 spectrometer (Agilent Technologies, Cork, Ireland). Quantitation  
 357 of each analyte was performed using MassHunter Quantitative  
 358 Analysis Software (Agilent Technologies, Cork, Ireland). The limit  
 359 of quantification was 1.32 pmol/g and 12.1 pmol/g, for AEA and  
 360 2-AG, respectively.

361 2.9. Stereotactic implantation of Guide Cannulae into the RVM

362 For experiment 3, stainless steel guide cannulae (Plastics One  
 363 Inc., Roanoke, VA, USA) were stereotactically implanted 1 mm  
 364 above the RVM (AP, 1.10 cm, ML  $\pm$  0.00 cm relative to bregma;  
 365 DV, 0.83 cm from dura) [66] under isoflurane anaesthesia (2%-3%  
 366 in O<sub>2</sub>; 0.5 L/min). The cannulae were permanently fixed to the skull  
 367 using stainless-steel screws and carboxylate cement. A stylet made  
 368 from stainless steel tubing (Plastics One Inc., Roanoke, VA) was  
 369 inserted into the guide cannulae to prevent blockage by debris.  
 370 The non-steroidal anti-inflammatory agent carprofen (5 mg/kg  
 371 s.c.) (Rimadyl; Pfizer, Kent, UK), and the broad-spectrum antibiotic  
 372 enrofloxacin (2.5 mg/kg s.c.) (Baytril; Bayer, Dublin, Ireland), were  
 373 administered before surgery to manage post-operative pain and to  
 374 prevent infection, respectively. After cannulae implantation, the  
 375 rats were housed singly and administered enrofloxacin (2.5 mg/  
 376 kg s.c.) for another 3 days. Rats were allowed to recover for at least  
 377 6 days before experimentation. During this period, the rats were  
 378 handled and their body weight and general health monitored on  
 379 a daily basis.

380 2.10. Histological verification of microinjection sites

381 For experiment 3, the sites of intra-cerebral microinjection  
 382 were determined before data analysis. Brain sections with fast-  
 383 green dye mark were collected on a cryostat (30- $\mu$ m thickness),  
 384 mounted on gelatinised glass slides, and counterstained with



**Fig. 2.** AEA (a) and 2-AG (b) levels in the RVM after intra-plantar saline or formalin administration in SD and WKY rats. Two-way analysis of variance for AEA (strain  $\times$  formalin interaction:  $F_{1,19} = 5.149, P = .037$ ) and 2-AG (strain  $\times$  formalin interaction:  $F_{1,23} = 7.148, P = .015$ ) followed by Fisher's least squares difference post-hoc test (\* $P = .023$  vs SD-Sal, \* $P = .047$  vs WKY-Sal, \* $P = .016$  and 0.023 vs SD-Form for AEA and 2-AG respectively). Data are expressed as mean  $\pm$  SEM ( $n = 5$  or 6). AEA, anandamide; 2-AG, 2-arachidonoyl glycerol; Form, formalin; Sal, saline solution; SD, Sprague-Dawley; WKY, Wistar-Kyoto.

385 cresyl violet to locate the precise position of microinjection sites  
386 under light microscopy.

387 2.11. Data analysis

388 The SPSS statistical package (IBM SPSS v20.0 for Windows; SPSS,  
389 Inc., Chicago, IL) was used to analyse all data. All data passed nor-  
390 mality testing (Shapiro–Wilk test). The time course of formalin-  
391 evoked nociceptive behaviour is presented in 5-minute bins for  
392 each study. Further analysis of data collapsed over extended peri-  
393 ods of the formalin trials or analysis of mRNA, neurochemical, or  
394 protein expression data was carried out using 2-way analysis of  
395 variance (ANOVA) followed by Fisher’s least squares difference  
396 (LSD) post hoc test where appropriate. Two-tailed unpaired  
397 Student *t* tests were performed to analyse neurochemical data from  
398 experiment 3. Data were considered significant at  $P < .05$ . Results  
399 are expressed as group mean  $\pm$  standard error of the mean (SEM).

400 3. Results

401 3.1. Formalin-evoked nociceptive responding is increased in WKY  
402 compared with SD counterparts


403 In Experiment 1, intra-plantar injection of formalin produced  
404 robust licking, biting, shaking, flinching and elevation of the

405 injected right hindpaw of both WKY and SD rats, expressed as  
406 the composite pain score (CPS). Formalin-treated rats of both  
407 strains displayed significantly greater nociceptive behaviour  
408 compared with saline-treated controls, in which composite pain  
409 scores were negligible (Fig. 1a SD/WKY-FORM vs SD/WKY-SAL,  
410  $P < .001$ ). However, WKY rats exhibited significantly greater  
411 formalin-evoked nociceptive behaviour compared with SD rats  
412 (Fig. 1b WKY-FORM vs SD-FORM,  $P < .001$ ) throughout the  
413 30-minute trial.

414 3.2. WKY rats exhibit a deficit in formalin-evoked mobilisation of AEA  
415 and 2-AG in the RVM

416 Data from mass spectrometry analysis of punch-dissected RVM  
417 tissue are presented in Fig 2.

418 Intra-plantar formalin administration to WKY rats was associ-  
419 ated with decreased levels of AEA, with no change in 2-AG levels  
420 (Fig. 2a WKY-Sal vs WKY-Form,  $P < .05$ ). In contrast, intra-plantar  
421 formalin administration to SD rats had no significant effect on  
422 AEA levels, but was associated with increased levels of 2-AG in  
423 the RVM (Fig. 2b SD-Sal vs SD-Form,  $P < .05$ ). Formalin-treated  
424 WKY rats had significantly lower levels of AEA and 2-AG when  
425 compared with SD counterparts (Fig. 2 SD-Form vs WKY-Form,  
426  $P < .05$ ).



**Fig. 3.** Intra-plantar formalin administration is associated with increased expression of NAPE-PLD and DAGLα mRNA in the RVM of SD, but not WKY, rats. Two-way analysis of variance for NAPE-PLD (formalin:  $F_{1,20} = 9.040$ ,  $P = .008$ ) and DAGL-α (formalin:  $F_{1,20} = 18.968$ ,  $P < .001$ , strain:  $F_{1,20} = 8.357$ ,  $P = .034$  and strain  $\times$  formalin interaction:  $F_{(1,20)} = 8.357$ ,  $P = .010$ ) mRNA levels followed by Fisher’s least squares difference post-hoc test (### $P < .001$  SD-Sal vs SD-Form; \* $P = 0.04$ , \*\* $P = 0.002$  SD-Form vs WKY-Form). All data are expressed as mean percentage of SD-saline control levels as mean  $\pm$  SEM ( $n = 5$  or  $6$ ). WKY, Wistar-Kyoto; DAGLα, diacylglycerol lipase α; FAAH, fatty acid amide hydrolase; Form, formalin; MAGL, monoacylglycerol lipase; NAPE-PLD, *N*-arachidonoyl-phosphatidylethanolamine phospholipase-D; Sal, saline solution; SD, Sprague-Dawley.

3.3. Intra-plantar formalin administration is associated with increased levels of NAPE-PLD and DAGL $\alpha$  mRNA in the RVM of SD, but not WKY, rats

There were no significant differences in levels of mRNA coding for the enzymes responsible for the synthesis (NAPE-PLD or DAGL $\alpha$ ) or catabolism (FAAH or MAGL) of AEA or 2-AG, or in levels of mRNA or protein for the CB $_1$  receptor, between saline-treated SD and WKY rats (Figs. 3 and 4, SD-Sal vs WKY-Sal). Intra-plantar formalin administration had no effect on levels of mRNA for MAGL, FAAH, or CB $_1$  receptor in either strain or on CB $_1$  receptor protein expression (Figs. 3 and 4, SD/WKY-Sal vs SD/WKY-Form). However, intra-plantar formalin administration was associated with increased levels of mRNA for NAPE-PLD (Fig. 3a, SD-Form vs SD-Sal,  $P = .01$ ) and DAGL $\alpha$  (Fig. 3b, SD-Form vs SD-Sal,  $P < .001$ ) in SD rats but not in WKY counterparts.

3.3.1. Increased formalin-evoked nociceptive responding in WKY rats is subject to modulation by pharmacological manipulation of the endocannabinoid system

In experiment 2, we repeated the finding of increased formalin-evoked nociceptive responding in WKY rats compared with SD rats, evident here over the first 40 minutes after formalin administration (Fig. 5b and c WKY-Veh vs SD-Veh,  $P < .001$ ). We focussed our efforts on pharmacological modulation of AEA rather than 2-AG because we observed a formalin-evoked reduction in the former, but not the latter, in the RVM of WKY rats. Systemic administration of the FAAH inhibitor URB597 (0.5 mg/kg i.p.), 60 minutes before formalin, significantly attenuated nociceptive behaviour in WKY rats over the first 40 minutes of the trial (Fig. 5b and c WKY-Veh vs WKY-URB597,  $P < .001$ ), whereas it had no effect in their SD counterparts (Fig. 5a and c SD-Veh vs SD-URB597). Conversely, systemic administration of the CB $_1$  receptor antagonist/inverse agonist AM251 (3 mg/kg), 30 minutes before formalin injection, significantly potentiated nociceptive responding in WKY rats 35–70 minutes after formalin administration (Fig. 5b and d, WKY-Veh vs WKY-AM251,  $P < 0.05$ ), while having no effect on formalin-evoked nociceptive behaviour in SD rats (Fig. 5a and d, SD-Veh vs SD-AM251).

3.4. Increased formalin-evoked nociceptive responding in WKY rats is subject to modulation by the endocannabinoid system in the RVM

Experiment 3 was carried out to evaluate whether the URB597-induced suppression of formalin-evoked hyperalgesia in the WKY rats, as observed in experiment 2, is mediated by AEA-induced activation of CB $_1$  receptors in the RVM. The reduction in formalin-evoked nociceptive behaviour after systemic administration of URB597 (0.5 mg/kg i.p.) to WKY rats that we observed in the first 35 minutes of the formalin trial in experiment 2 was apparent over the first 15 minutes in experiment 3 (Fig. 6a and b VEH-DMSO vs URB597-DMSO,  $P < .05$ ), and was prevented by microinjection of the CB $_1$  receptor antagonist AM251 (1.0  $\mu$ g/0.3  $\mu$ l) directly into the RVM (Fig. 6a and b, URB597-DMSO vs URB597-AM251,  $P < .05$ ). In all, 80% of the intracerebral microinjections were placed within the borders of the RVM, with the remaining injections positioned proximal to, but outside the borders of, this region (Fig. 8). Only the results of experiments in which microinjections were accurately positioned within the borders of the RVM were included in the analyses.

3.5. Pharmacological effect of URB597 administration on AEA and 2-AG levels in the RVM of saline- or formalin-treated SD and WKY rats

Measurement of endocannabinoid levels in the RVM of WKY rats at the end of experiment 3 revealed that URB597 significantly





Fig. 4. Expression of (a) glycosylated (~62 kDa) or (b) non-glycosylated (~53 kDa) CB $_1$  receptor in RVM tissue from saline- or formalin-injected SD and WKY rats. Two-way ANOVA revealed no significant effect of strain, formalin administration or their interaction. Data are expressed as mean percentage of SD-saline control  $\pm$  SEM ( $n = 6$ ). Form, formalin; RVM, rostromedial medulla; SD, Sal, saline solution; Sprague–Dawley; WKY, Wistar–Kyoto.


increased levels of AEA, but not 2-AG, in the RVM of WKY rats (Fig. 7 WKY-Veh vs WKY-URB597;  $t_8 = 2.464$ ,  $P < .05$  for AEA and  $t_8 = 0.980$ ,  $P = 0.36$  for 2-AG).

#### 4. Discussion


The data presented here suggest that impaired endocannabinoid-CB $_1$  receptor signalling in the RVM underpins the hyperalgesic response of stress-hyperresponsive WKY rats to intra-plantar formalin injection, compared with stress-normoresponsive SD rats. The magnitude of formalin-evoked nociceptive behaviour in WKY rats was greater than in SD rats, and this hyperalgesic phenotype was attenuated by inhibition of AEA catabolism and exacerbated by CB $_1$  receptor blockade. Moreover, the increased formalin-evoked nociceptive response in WKY rats was associated with lower tissue levels of AEA and 2-AG, and blunted formalin-evoked



**Fig. 5.** AM251 (3 mg/kg i.p.) potentiates, whereas URB597 (0.5 mg/kg i.p.) attenuates, the enhanced nociceptive response of WKY rats to formalin administration. (a and b) Temporal profile of formalin-evoked nociceptive behaviour in SD and WKY rats after AM251 and URB597 administration. (c and d) Graphic representations of collapsed data analysed over periods indicated by boxed areas in panels a and b. Two-way analysis of variance (a) 0 to 40 minutes after formalin administration (strain:  $F_{1,62} = 14.650$ ,  $P < 0.001$ ; drug:  $F_{2,62} = 16.899$ ,  $P < .001$  and strain  $\times$  drug interaction:  $F_{2,62} = 8.685$ ,  $P = .001$ ) and (b) 35 to 70 minutes after formalin (drug:  $F_{2,62} = 3.324$ ,  $P = .043$ ), followed by Fisher's least squares difference post-hoc test (\*\* $P < .01$  vs SD-Veh, \* $P = .039$ , \*\*\* $P < .001$  vs WKY-Veh and # $P = .032$ , ### $P < .001$  vs SD-AM251). Data are expressed as mean  $\pm$  SEM ( $n = 10-12$ ); SD, Sprague-Dawley; Veh, vehicle; WKY, Wistar-Kyoto;.



**Fig. 6.** Microinjection of AM251 (1.0  $\mu$ g/0.3  $\mu$ L) directly into the RVM of WKY rats prevented the systemic URB597 (0.5 mg/kg i.p.)-induced reduction in formalin-evoked nociceptive behaviour over the first 15 minutes of the 70-minute trial. (a) Temporal profile of formalin-evoked nociceptive behaviour in cannulated WKY rats. (b) Graphic representations of collapsed data analysed over the period indicated by the boxed area in panel a. Two-way analysis of variance (systemic drug  $\times$  intra-RVM drug interaction:  $F_{1,23} = 9.214$ ,  $P = .007$ ) followed by Fisher's LSD post-hoc test (\* $P = .012$  vs VEH-DMSO and # $P = 0.014$  vs URB597-DMSO). Data are expressed as mean  $\pm$  SEM ( $n = 5-8$ ).




**Fig. 7.** Systemic URB597 (0.5 mg/kg i.p.) administration significantly increases AEA levels in the RVM of formalin-treated WKY rats receiving DMSO microinjection into the RVM. Two-tailed  $t$  test for AEA ( $t_8 = 2.464$ ,  $P = .039$ ) and 2-AG ( $t_8 = 0.980$ ,  $P = .356$ ) levels. Data are expressed as mean  $\pm$  SEM ( $n = 5$ ). AEA, anandamide; 2-AG, 2-arachidonoyl glycerol; WKY, Wistar-Kyoto; DMSO, dimethyl sulfoxide; RVM, rostroventromedial medulla.

501 induction of NAPE-PLD and DAGL $\alpha$  mRNA, in the RVM, compared with those in SD controls. Furthermore, pharmacological blockade  
502

of CB $_1$  receptors in the RVM attenuated the antinociceptive effect of the FAAH inhibitor URB597 in WKY rats.

503  
504



**Fig. 8.** Diagrammatic representation of the confirmed sites of microinjection of dimethyl sulfoxide (DMSO) (●) or AM251 (■) into the rostroventromedial medulla of Wistar-Kyoto rats receiving systemic administration of (a) vehicle or (b) URB597. Distances are relative to bregma. (Based on Paxinos and Watson [66]).

In this study, we replicated our earlier finding of increased nociceptive behaviour in WKY rats as compared to the SD comparator strain over the first 35 minutes of the formalin test [5]. We sought to extend these findings by investigating potential neurochemical and receptor mechanisms underlying the hyperalgesic phenotype, with a focus on the endocannabinoid system in the RVM given its key role in top-down descending modulation of pain [21,22,91] and evidence that CB<sub>1</sub> receptors in the RVM regulate nociceptive processing [36,50,55,84]. Levels of AEA and 2-AG, mRNA coding for catabolizing or synthesizing enzymes or CB<sub>1</sub> receptor mRNA or protein expression were similar in the RVM of SD and WKY rats receiving an intra-plantar injection of saline, suggesting that the resting endocannabinoid tone in the RVM is similar between the 2 strains. Formalin-evoked nociceptive behaviour was associated with increased levels of 2-AG in SD rats, and decreased AEA in the RVM of WKY rats. When compared with formalin-treated SD counterparts, RVM tissue levels of both AEA and 2-AG were lower in formalin-treated WKY rats. In addition, formalin-evoked nociceptive behaviour in SD rats was associated with increased RVM tissue levels of mRNA coding for NAPE-PLD and DAGL- $\alpha$ , the enzymes responsible for the synthesis of AEA and 2-AG, respectively. No such formalin-evoked increases were observed in WKY rats. Formalin injection had no effects on mRNA levels for the endocannabinoid catabolizing enzymes or on CB<sub>1</sub> receptor mRNA or protein expression in the RVM of either strain. These data suggest differential formalin-evoked recruitment of the endocannabinoid system in the RVM in 2 rat strains differing in their responsivity to an inflammatory noxious stimulus. Overall, our findings indicate impaired mobilization of endocannabinoids, and suppressed mRNA expression of genes coding for enzymes that synthesise the endocannabinoids, in the RVM of WKY rats in response to intra-plantar formalin injection. These alterations suggest reduced endocannabinoid tone in the RVM of WKY rats in response to a noxious insult, and we hypothesised that these alterations may play an important role in the hyperalgesic phenotype of WKY rats.

Subsequent pharmacological manipulation studies confirmed a role for the endocannabinoid system in the hyperalgesic phenotype expressed in WKY rats. We focussed our efforts on pharmacological modulation of AEA rather than 2-AG because we observed a formalin-evoked reduction in the former, but not the latter, in the RVM of WKY rats. Systemic administration of the FAAH inhibitor URB597

reduced, whereas the CB<sub>1</sub> receptor antagonist AM251 potentiated, formalin-evoked nociceptive behaviour in WKY rats. In contrast, neither drug had any effect on formalin-evoked nociceptive behaviour in SD rats at the doses administered here. We propose that a deficit in formalin-evoked endocannabinoid tone in the CNS of WKY rats may explain the efficacy of these 2 endocannabinoid system modulators in WKY, but not SD, rats; that is, against a background of reduced endocannabinoid tone in discrete brain regions such as the RVM, as indicated from our reported tissue levels of AEA and 2-AG from experiment 1, pharmacological inhibition of FAAH and blockade of CB<sub>1</sub> may be more effective in modulating pain-related behaviour in WKY rats than in the SD strain, in which endocannabinoid tone was greater. Thus, in WKY rats, FAAH inhibition would be expected to restore endocannabinoid tone and reduce hyperalgesia, whereas CB<sub>1</sub> receptor blockade would further exacerbate the deficit in endocannabinoid tone, thereby potentiating hyperalgesia, as was observed. Our study is the first to investigate the effects of these endocannabinoid system modulators on nociceptive behaviour in WKY rats. Furthermore, our data support earlier findings in which URB597 [34] and AM251 [1,9] failed to alter formalin-evoked nociceptive responses in SD rats. In comparison, the CB<sub>1</sub> receptor antagonist rimonabant has been reported to modestly increase formalin-evoked nociceptive behaviour in SD rats when administered 5 minutes after formalin [82].

Further direct evidence for a role of the endocannabinoid system in the RVM in the modulation of pain is provided by drug microinjection and electrophysiological studies. Microinjection of cannabinoid compounds directly into the RVM have been shown to modulate neuronal firing in cells of the RVM under conditions of acute [50,54,55,57] and persistent [57] inflammatory pain, as determined by in vivo electrophysiological studies. The RVM has a dense population of serotonergic [3,11,46,61,78], GABAergic [2,56], and glutamatergic neurons [81]; however, the expression of CB<sub>1</sub> receptors on GABAergic, glutamatergic, or serotonergic neurons in the RVM has yet to be confirmed anatomically (for review, see Rea et al. [70]). Moreover, to our knowledge, no studies to date have investigated serotonergic, GABAergic, or glutamatergic tone in the RVM of WKY vs SD rats. As such, the precise neurochemical mechanisms by which endocannabinoids in the RVM influence hyperalgesia in the WKY rat requires further scrutiny and should be the subject of future studies. In an animal model of unconditioned

stress-induced analgesia (SIA) involving exposure to footshock with subsequent assessment of rat tail-flick responses, microinjection of the CB<sub>1</sub> receptor antagonist rimonabant into the RVM suppressed SIA, whereas intra-RVM administration of URB597 enhanced SIA [84]. In the present study, we used site-specific drug microinjection methodology to further investigate the role of CB<sub>1</sub> receptors in the RVM in mediating the antinociceptive effects of systemically administered URB597 in WKY rats, and investigated changes in endocannabinoid levels associated with URB597 administration. In RVM-cannulated WKY rats, systemic administration of URB597 reduced formalin-evoked nociceptive behaviour, albeit with an onset and duration of action that were earlier and shorter, respectively, than was observed in the WKY rats in experiment 2 that were not cannulated and did not receive intra-RVM DMSO. The suppression of formalin-evoked nociceptive behaviour in WKY rats by the FAAH inhibitor URB597 was associated with increased levels of AEA, but not 2-AG, in the RVM. Intra-RVM administration of AM251 prevented the URB597-induced reduction in formalin-evoked nociceptive behaviour, whereas it had no effect on nociceptive behaviour when administered alone. Taken with the data discussed above, these results together strongly suggest that the URB597-mediated reduction in formalin-evoked nociceptive behaviour in WKY rats is mediated through the activation of CB<sub>1</sub> receptors in the RVM by elevated levels of AEA arising from FAAH inhibition. In addition to activating CB<sub>1</sub> receptors, AEA is also an agonist at the transient receptor potential vanilloid subtype 1 (TRPV1) channel [13,16,74,79], the expression of which has been reported in the RVM [80]. However, to our knowledge, there are no studies investigating the effects of administration of TRPV1 agonists or antagonists directly into the RVM on nociceptive behaviour, and so it is uncertain whether TRPV1 in the RVM plays a direct role in the regulation of nociception. However, given that the antinociceptive effects of systemically administered URB597 were blocked completely by intra-RVM administration of the selective CB<sub>1</sub> receptor antagonist AM251 in the present study, and given that AM251 does not have any activity directly at TRPV1 [65], it seems very likely that the antinociceptive effects of URB597 in the present study are mediated exclusively by CB<sub>1</sub> receptors, without any direct involvement of TRPV1. Although we cannot definitively rule out the possibility that blockade of the CB<sub>1</sub> receptor activates non-CB<sub>1</sub> targets (eg, GPR55) or shunts AEA towards activation of other receptors in the RVM, it also seems unlikely that such mechanisms can explain the effects observed herein, because (1) intra-RVM administration of AM251 alone had no effect on formalin-evoked nociceptive behaviour, and (2) URB597 (and the resulting AEA elevation) had no effect on formalin-evoked nociceptive behaviour in animals receiving intra-RVM AM251. Overall, these data support the contention that hyperalgesia in the WKY rats is mediated, at least in part, via a deficit in endocannabinoid-CB<sub>1</sub> tone in the RVM.

#### 4.1. Conclusions

In conclusion, our data provide evidence for an altered nociceptive response in a genetic strain predisposed to negative affect. The results indicate a compromised endocannabinoid system in the RVM of WKY rats compared with SD controls, a key neuroanatomical brain region involved in descending pain modulation. Dysfunction of the endocannabinoid system in this top-down control system may be maladaptive, contributing to exacerbated responsiveness to noxious stimuli. Pharmacological normalisation of this endocannabinoid system dysfunction attenuated this genotype-dependent hyperalgesia in rodents and may represent a useful and novel therapeutic approach for the treatment of patients with pain that is exacerbated by negative affect or co-morbid with stress-related psychiatric disorders.

#### Conflict of interest statement

The authors have no conflicts of interest to declare.

#### Acknowledgements

This work was supported by a research grant from Science Foundation Ireland (10/IN.1/B2976).

#### References

- [1] Borsani E, Labanca M, Bianchi R, Rodella LF. AM404 decreases Fos-immunoreactivity in the spinal cord in a model of inflammatory pain. *Brain Res* 2007;1152:87–94.
- [2] Bowery NG, Hudson AL, Price GW. GABAA and GABAB receptor site distribution in the rat central nervous system. *Neuroscience* 1987;20:365–83.
- [3] Bowker RM, Westlund KN, Coulter JD. Origins of serotonergic projections to the spinal cord in rat: an immunocytochemical-retrograde transport study. *Brain Res* 1981;226:187–99.
- [4] Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–54.
- [5] Burke NN, Hayes E, Calpin P, Kerr DM, Moriarty O, Finn DP, Roche M. Enhanced nociceptive responding in two rat models of depression is associated with alterations in monoamine levels in discrete brain regions. *Neuroscience* 2010;171:1300–13.
- [6] Butler RK, Finn DP. Stress-induced analgesia. *Prog Neurobiol* 2009;88:184–202.
- [7] Butler RK, Ford GK, Hogan M, Roche M, Doyle KM, Kelly JP, Kendall DA, Chapman V, Finn DP. Fear-induced suppression of nociceptive behaviour and activation of Akt signalling in the rat periaqueductal grey: role of fatty acid amide hydrolase. *J Psychopharmacol* 2012;26:83–91.
- [8] Butler RK, Rea K, Lang Y, Gavin AM, Finn DP. Endocannabinoid-mediated enhancement of fear-conditioned analgesia in rats: opioid receptor dependency and molecular correlates. *PAIN<sup>®</sup>* 2008;140:491–500.
- [9] Cornelio AM, Nunes-de-Souza RL, Morgan MM. Contribution of the rostral ventromedial medulla to post-anxiety induced hyperalgesia. *Brain Res* 2012;1450:80–6.
- [10] Cravatt BF, Giang DK, Mayfield SP, Boger DL, Lerner RA, Gilula NB. Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature* 1996;384:83–7.
- [11] Dahlstrom A, Fuxe K. Localization of monoamines in the lower brain stem. *Experientia* 1964;20:398–9.
- [12] de Novellis V, Palazzo E, Rossi F, De Petrocellis L, Petrosino S, Guida F, Luongo L, Migliozi A, Cristino L, Marabese I, Starowicz K, Di Marzo V, Maione S. The analgesic effect of N-arachidonoyl-serotonin, a FAAH inhibitor and TRPV1 receptor antagonist, associated with changes in rostral ventromedial medulla and locus coeruleus cell activity in rats. *Neuropharmacology* 2008;55:1105–13.
- [13] De Petrocellis L, Bisogno T, Maccarrone M, Davis JB, Finazzi-Agrilli A, Di Marzo V. The activity of anandamide at vanilloid VR1 receptors requires facilitated transport across the cell membrane and is limited by intracellular metabolism. *J Biol Chem* 2001;276:12856–63.
- [14] Devane WA, Dysarz 3rd FA, Johnson MR, Melvin LS, Howlett AC. Determination and characterization of a cannabinoid receptor in rat brain. *Mol Pharmacol* 1988;34:605–13.
- [15] Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, Griffin G, Gibson D, Mandelbaum A, Etinger A, Mechoulam R. Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* 1992;258:1946–9.
- [16] Di Marzo V, Bisogno T, De Petrocellis L. Anandamide: some like it hot. *Trends Pharmacol Sci* 2001;22:346–9.
- [17] Di Marzo V, Petrosino S. Endocannabinoids and the regulation of their levels in health and disease. *Curr Opin Lipidol* 2007;18:129–40.
- [18] Ebrahimzadeh M, Haghparast A. Analgesic effects of cannabinoid receptor agonist WIN5,212–2 in the nucleus cuneiformis in animal models of acute and inflammatory pain in rats. *Brain Res* 2011;1420:19–28.
- [19] Egertova M, Elphick MR. Localisation of cannabinoid receptors in the rat brain using antibodies to the intracellular C-terminal tail of CB<sub>1</sub>. *J Comp Neurol* 2000;422:159–71.
- [20] Escobar W, Ramirez K, Avila C, Limongi R, Vanegas H, Vazquez E. Metamizol, a non-opioid analgesic, acts via endocannabinoids in the PAG-RVM axis during inflammation in rats. *Eur J Pain* 2012;16:676–89.
- [21] Fields HL, Basbaum AI. Brainstem control of spinal pain-transmission neurons. *Annu Rev Physiol* 1978;40:217–48.
- [22] Fields HL, Heinricher MM, Mason P. Neurotransmitters in nociceptive modulatory circuits. *Annu Rev Neurosci* 1991;14:219–45.
- [23] Finn DP. Endocannabinoid-mediated modulation of stress responses: physiological and pathophysiological significance. *Immunobiology* 2009;215:629–46.
- [24] Finn DP, Beckett SR, Richardson D, Kendall DA, Marsden CA, Chapman V. Evidence for differential modulation of conditioned aversion and fear-conditioned analgesia by CB<sub>1</sub> receptors. *Eur J Neurosci* 2004;20:848–52.

- [25] Finn DP, Chapman V, Jhaveri MD, Samanta S, Manders T, Bowden J, Matthews L, Marsden CA, Beckett SR. The role of the central nucleus of the amygdala in nociception and aversion. *Neuroreport* 2003;14:981–4.
- [26] Finn DP, Jhaveri MD, Beckett SR, Madjd A, Kendall DA, Marsden CA, Chapman V. Behavioral, central monoaminergic and hypothalamo–pituitary–adrenal axis correlates of fear-conditioned analgesia in rats. *Neuroscience* 2006;138:1309–17.
- [27] Finn DP, Viveros MP, Marco EM. The endocannabinoid system and emotional processing: Pathophysiology and therapeutic potential. *J Psychopharmacol* 2012;26:3–6.
- [28] Ford GK, Kieran S, Dolan K, Harhen B, Finn DP. A role for the ventral hippocampal endocannabinoid system in fear-conditioned analgesia and fear responding in the presence of nociceptive tone in rats. *PAIN<sup>®</sup>* 2011;152:2495–504.
- [29] Gentsch C, Lichtsteiner M, Feer H. Genetic and environmental influences on behavioral and neurochemical aspects of emotionality in rats. *Experientia* 1988;44:482–90.
- [30] Giang DK, Cravatt BF. Molecular characterization of human and mouse fatty acid amide hydrolases. *Proc Natl Acad Sci* 1997;94:2238–42.
- [31] Gibney SM, Gosselin RD, Dinan TG, Cryan JF. Colorectal distension-induced prefrontal cortex activation in the Wistar-Kyoto rat: Implications for irritable bowel syndrome. *Neuroscience* 2010;165:675–83.
- [32] Gonsiorek W, Lunn C, Fan X, Narula S, Lundell D, Hipkin RW. Endocannabinoid 2-arachidonyl glycerol is a full agonist through human type 2 cannabinoid receptor: Antagonism by anandamide. *Mol Pharmacol* 2000;57:1045–50.
- [33] Gunter WD, Shepard JD, Foreman RD, Myers DA, Greenwood-Van Meerveld B. Evidence for visceral hypersensitivity in high-anxiety rats. *Physiol Behav* 2000;69:379–82.
- [34] Hasanein P, Parviz M, Keshavarz M, Roohbakhsh A. URB597, an inhibitor of fatty acid amide hydrolase, reduces hyperalgesia in diabetic rats. *Can J Physiol Pharmacol* 2009;87:432–9.
- [35] Hasanein P, Shahidi S, Komaki A, Mirazi N. Effects of URB597 as an inhibitor of fatty acid amide hydrolase on modulation of nociception in a rat model of cholestasis. *Eur J Pharmacol* 2008;591:132–5.
- [36] Heinricher MM, Tavares I, Leith JL, Lumb BM. Descending control of nociception: specificity, recruitment and plasticity. *Brain Res Rev* 2009;60:214–25.
- [37] Herkenham M, Lynn AB, Johnson MR, Melvin LS, de Costa BR, Rice KC. Characterization and localization of cannabinoid receptors in rat brain: a quantitative in vitro autoradiographic study. *J Neurosci* 1991;11:563–83.
- [38] Hill MN, McLaughlin RJ, Bingham B, Shrestha L, Lee TT, Gray JM, Hillard CJ, Gorzalka BB, Viau V. Endogenous cannabinoid signaling is essential for stress adaptation. *Proc Natl Acad Sci U S A* 2010;107:9406–11.
- [39] Hohmann AG, Suplita 2nd RL. Endocannabinoid mechanisms of pain modulation. *AAPS J* 2006;8:E693–708.
- [40] Hong S, Fan J, Kemmerer ES, Evans S, Li Y, Wiley JW. Reciprocal changes in vanilloid (TRPV1) and endocannabinoid (CB1) receptors contribute to visceral hyperalgesia in the water avoidance stressed rat. *Gut* 2009;58:202–10.
- [41] Jayamanne A, Greenwood R, Mitchell VA, Aslan S, Piomelli D, Vaughan CW. Actions of the FAAH inhibitor URB597 in neuropathic and inflammatory chronic pain models. *Br J Pharmacol* 2006;147:281–8.
- [42] Kerr DM, Burke NN, Ford GK, Connor TJ, Harhen B, Egan LJ, Finn DP, Roche M. Pharmacological inhibition of endocannabinoid degradation modulates the expression of inflammatory mediators in the hypothalamus following an immunological stressor. *Neuroscience* 2012;204:53–63.
- [43] Kerr DM, Harhan B, Okine BN, Egan LJ, Finn DP, Roche M. The monoacylglycerol lipase inhibitor JZL184 attenuates LPS-induced increases in cytokine expression in the rat frontal cortex and plasma: Differential mechanisms of action. *Br J Pharmacol* 2013;169:808–19.
- [44] Kinsey SG, Long JZ, O'Neal ST, Abdullah RA, Poklis JL, Boger DL, Cravatt BF, Lichtman AH. Blockade of endocannabinoid-degrading enzymes attenuates neuropathic pain. *J Pharmacol Exp Ther* 2009;330:902–10.
- [45] Liu J, Wang L, Harvey-White J, Huang BX, Kim H-Y, Luquet S, Palmiter RD, Krystal G, Rai R, Mahadevan A, Razdan RK, Kunos G. Multiple pathways involved in the biosynthesis of anandamide. *Neuropharmacology* 2008;54:1–7.
- [46] Lovick TA, Robinson JP. Bulbar raphe neurones with projections to the trigeminal nucleus caudalis and the lumbar cord in the rat: A fluorescence double-labelling study. *Exp Brain Res* 1983;50:299–308.
- [47] Luk T, Jin W, Zvonok A, Lu D, Lin XZ, Chavkin C, Makriyannis A, Mackie K. Identification of a potent and highly efficacious, yet slowly desensitizing CB1 cannabinoid receptor agonist. *Br J Pharmacol* 2004;142:495–500.
- [48] Lutz B. Endocannabinoid signals in the control of emotion. *Curr Opin Pharmacol* 2009;9:46–52.
- [49] Maione S, Bisogno T, de Novellis V, Palazzo E, Cristino L, Valenti M, Petrosino S, Guglielmotti V, Rossi F, Di Marzo V. Elevation of endocannabinoid levels in the ventrolateral periaqueductal grey through inhibition of fatty acid amide hydrolase affects descending nociceptive pathways via both cannabinoid receptor type 1 and transient receptor potential vanilloid type-1 receptors. *J Pharmacol Exp Ther* 2006;316:969–82.
- [50] Martin WJ, Tsou K, Walker JM. Cannabinoid receptor-mediated inhibition of the rat tail-flick reflex after microinjection into the rostral ventromedial medulla. *Neurosci Lett* 1998;242:33–6.
- [51] Matsuda LA, Bonner TI, Lolait SJ. Localization of cannabinoid receptor mRNA in rat brain. *J Comp Neurol* 1993;327:535–50.
- [52] Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 1990;346:561–4.
- [53] Mechoulam R, Ben-Shabat S, Hanus L, Ligumsky M, Kaminski NE, Schatz AR, Gopher A, Almog S, Martin BR, Compton DR, Pertwee RG, Griffin G, Bayewitch M, Barg J, Vogel Z. Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. *Biochem Pharmacol* 1995;50:83–90.
- [54] Meng ID, Johansen JP. Antinociception and modulation of rostral ventromedial medulla neuronal activity by local microinfusion of a cannabinoid receptor agonist. *Neuroscience* 2004;124:685–93.
- [55] Meng ID, Manning BH, Martin WJ, Fields HL. An analgesia circuit activated by cannabinoids. *Nature* 1998;395:381–3.
- [56] Millhorn DE, Hokfelt T, Seroogy K, Verhofstad AA. Extent of colocalization of serotonin and GABA in neurons of the ventral medulla oblongata in rat. *Brain Res* 1988;461:169–74.
- [57] Monhemius R, Azami J, Green DL, Roberts MH. CB1 receptor mediated analgesia from the nucleus reticularis gigantocellularis pars alpha is activated in an animal model of neuropathic pain. *Brain Res* 2001;908:67–74.
- [58] Munro S, Thomas KL, Abu-Shaar M. Molecular characterization of a peripheral receptor for cannabinoids. *Nature* 1993;365:61–5.
- [59] Okine B, Norris L, Woodhams S, Burston J, Patel A, Alexander S, Barrett D, Kendall D, Bennett A, Chapman V. Lack of effect of chronic pre-treatment with the FAAH inhibitor URB597 on inflammatory pain behaviour: Evidence for plastic changes in the endocannabinoid system. *Br J Pharmacol* 2012;167:627–40.
- [60] Olango WM, Roche M, Ford GK, Harhen B, Finn DP. The endocannabinoid system in the rat dorsolateral periaqueductal grey mediates fear-conditioned analgesia and controls fear expression in the presence of nociceptive tone. *Br J Pharmacol* 2011;165:2549–60.
- [61] Oliveras JL, Bourgoin S, Hery F, Besson JM, Hamon M. The topographical distribution of serotonergic terminals in the spinal cord of the cat: biochemical mapping by the combined use of microdissection and microassay procedures. *Brain Res* 1977;138:393–406.
- [62] Otrubova K, Ezzili C, Boger DL. The discovery and development of inhibitors of fatty acid amide hydrolase (FAAH). *Bioorgan Med Chem Lett* 2011;21:4674–85.
- [63] Pardon MC, Gould GG, Garcia A, Phillips L, Cook MC, Miller SA, Mason PA, Morilak DA. Stress reactivity of the brain noradrenergic system in three rat strains differing in their neuroendocrine and behavioral responses to stress: Implications for susceptibility to stress-related neuropsychiatric disorders. *Neuroscience* 2002;115:229–42.
- [64] Pare WP. The performance of WKY rats on three tests of emotional behavior. *Physiol Behav* 1992;51:1051–6.
- [65] Patil M, Patwardhan A, Salas MM, Hargreaves KM, Akopian AN. Cannabinoid receptor antagonists AM251 and AM630 activate TRPA1 in sensory neurons. *Neuropharmacology* 2011;61:778–88.
- [66] Paxinos G, Watson C. The rat brain in stereotaxic coordinates. Academic Press; 1997.
- [67] Rea K, Lang Y, Finn DP. Alterations in extracellular levels of gamma-aminobutyric acid in the rat basolateral amygdala and periaqueductal gray during conditioned fear, persistent pain and fear-conditioned analgesia. *J Pain* 2009;10:1088–98.
- [68] Rea K, Olango WM, Harhen B, Kerr DM, Golligan R, Fitzgerald S, Moore M, Roche M, Finn DP. Evidence for a role of GABAergic and glutamatergic signalling in the basolateral amygdala in endocannabinoid-mediated fear-conditioned analgesia in rats. *PAIN<sup>®</sup>* 2013;154:576–85.
- [69] Rea K, Roche M, Finn DP. Modulation of conditioned fear, fear-conditioned analgesia, and brain regional c-Fos expression following administration of muscimol into the rat basolateral amygdala. *J Pain* 2011;12:712–21.
- [70] Rea K, Roche M, Finn DP. Supraspinal modulation of pain by cannabinoids: The role of GABA and glutamate. *Br J Pharmacol* 2007;152:633–48.
- [71] Robbins M, DeBerry J, Ness T. Chronic psychological stress enhances nociceptive processing in the urinary bladder in high-anxiety rats. *Physiol Behav* 2007;91:544–50.
- [72] Roche M, Johnston P, Mhuircheartaigh ON, Olango WM, Mackie K, Finn DP. Effects of intra-basolateral amygdala administration of rimonabant on nociceptive behaviour and neuronal activity in the presence or absence of contextual fear. *Eur J Pain* 2010;14:487–95.
- [73] Roche M, O'Connor E, Diskin C, Finn DP. The effect of CB(1) receptor antagonism in the right basolateral amygdala on conditioned fear and associated analgesia in rats. *Eur J Neurosci* 2007;26:2643–53.
- [74] Ross RA, Gibson TM, Brockie HC, Leslie M, Pashmi G, Craib SJ, Di Marzo V, Pertwee RG. Structure-activity relationship for the endogenous cannabinoid, anandamide, and certain of its analogues at vanilloid receptors in transfected cells and vas deferens. *Br J Pharmacol* 2001;132:631–40.
- [75] Schlosburg JE, Kinsey SG, Lichtman AH. Targeting fatty acid amide hydrolase (FAAH) to treat pain and inflammation. *AAPS J* 2009;11:39–44.
- [76] Seierstad M, Breitenbucher JG. Discovery and development of fatty acid amide hydrolase (FAAH) inhibitors. *J Medic Chem* 2008;51:7327–43.
- [77] Shen L, Yang XJ, Qian W, Hou XH. The role of peripheral cannabinoid receptors type 1 in rats with visceral hypersensitivity induced by chronic restraint stress. *J Neurogastroenterol Motil* 2010;16:281–90.

- 897 [78] Skagerberg G, Bjorklund A. Topographic principles in the spinal projections of  
898 serotonergic and non-serotonergic brainstem neurons in the rat. *Neuroscience*  
899 1985;15:445–80. 918
- 900 [79] Smart D, Gunthorpe MJ, Jerman JC, Nasir S, Gray J, Muir AI, Chambers JK,  
901 Randall AD, Davis JB. The endogenous lipid anandamide is a full agonist at the  
902 human vanilloid receptor (hVR1). *Br J Pharmacol* 2000;129:227–30. 919
- 903 [80] Starowicz K, Maione S, Cristino L, Palazzo E, Marabese I, Rossi F, de Novellis V,  
904 Di Marzo V. Tonic endovanilloid facilitation of glutamate release in brainstem  
905 descending antinociceptive pathways. *J Neurosci* 2007;27:13739–49. 920
- 906 [81] Starowicz K, Makuch W, Osikowicz M, Piscitelli F, Petrosino S, Di Marzo V,  
907 Przewlocka B. Spinal anandamide produces analgesia in neuropathic rats:  
908 Possible CB(1)- and TRPV1-mediated mechanisms. *Neuropharmacology* 2011;62:1746–55. 921
- 909 [82] Strangman NM, Patrick SL, Hohmann AG, Tsou K, Walker JM. Evidence for a  
910 role of endogenous cannabinoids in the modulation of acute and tonic pain  
911 sensitivity. *Brain Res* 1998;813:323–8. 922
- 912 [83] Sugiura T, Kondo S, Sukagawa A, Nakane S, Sindo A, Itoh K, Yamashita A, Waku  
913 K. 2-Arachidonoylglycerol: a possible endogenous cannabinoid receptor ligand  
914 in brain. *Biochem Biophys Res Commun* 1995;215:89–97. 923
- 915 [84] Suplita 2nd RL, Farthing JN, Gutierrez T, Hohmann AG. Inhibition of fatty-acid  
916 amide hydrolase enhances cannabinoid stress-induced analgesia: Sites of  
917 action in the dorsolateral periaqueductal gray and rostral ventromedial  
918 medulla. *Neuropharmacology* 2005;49:1201–9. 919
- [85] Taylor BK, Roderick RE, St Lezin E, Basbaum AI. Hypoalgesia and hyperalgesia  
920 with inherited hypertension in the rat. *Am J Physiol Regul Integr Comp Physiol*  
921 2001;280:R345–54. 922
- [86] Vinod KY, Xie S, Psychoyos D, Hungund BL, Cooper TB, Tejani-Butt SM.  
923 Dysfunction in fatty acid amide hydrolase is associated with depressive-like  
924 behavior in Wistar Kyoto rats. *PLoS One* 2012;7:e36743. 925
- [87] Viveros MP, Marco EM, File SE. Endocannabinoid system and stress and  
926 anxiety responses. *Pharmacol Biochem Behav* 2005;81:331–42. 927
- [88] Viveros MP, Marco EM, Llorente R, Lopez-Gallardo M. Endocannabinoid  
928 system and synaptic plasticity: implications for emotional responses. *Neural*  
929 *Plast* 2007;52908. 930
- [89] Watson GS, Sufka KJ, Coderre TJ. Optimal scoring strategies and weights for  
931 the formalin test in rats. *PAIN<sup>®</sup>* 1997;70:53–8. 932
- [90] Zeng Q, Wang S, Lim G, Yang L, Mao J, Sung B, Chang Y, Lim JA, Guo G.  
933 Exacerbated mechanical allodynia in rats with depression-like behavior.  
934 *Brain Res* 2008;1200:27–38. 935
- [91] Zhuo M, Gebhart GF. Biphasic modulation of spinal nociceptive transmission  
936 from the medullary raphe nuclei in the rat. *J Neurophysiol* 1997;78:746–58. 937
- 938

UNCORRECTED PROOF